arXiv:1107.1741v2 [math.OA] 17 Nov 2011

K-HOMOLOGY AND INDEX THEORY ON CONTACT MANIFOLDS

PAUL F. BAUM AND ERIK VAN ERP

Many of the observations in this paper one way or another go back to basic ideas of M. F. Atiyah. With admiration and affection we dedicate this paper to Sir Michael.

Contents

1. Introduction 2
1.1. Context 2
1.2. The problem 3
1.3. Statement of the result 6
1.4. Outline of the proof 9
1.5. Summary of the paper 12
Acknowledgment 12
2. The Poincaré duality map b 12
2.1. The C^*-algebra of the Heisenberg group 13
2.2. $U(n)$ symmetry 14
2.3. Bargmann-Fok space 15
2.4. Quantization 16
2.5. Associated bundles 17
2.6. Inverting the Connes-Thom isomorphism 18
2.7. Poincaré duality 20
3. The index theorem as a commutative triangle 22
3.1. Differential operators in the Heisenberg calculus 22
3.2. Heisenberg pseudodifferential calculus 23
3.3. The Heisenberg symbol in K-theory 25
3.4. The "Choose an operator" maps 26
3.5. The hypoelliptic index theorem in K-homology 28
4. Computation of the K-cycle $b(\sigma_H(P))$ 29
4.1. Perturbing the symbol 29
4.2. Suspension in K-theory 30
4.3. Second order scalar operators 31
4.4. Toeplitz operators 34
4.5. The general case: vector bundles 35
4.6. A hypoelliptic index problem posed by R. Ponge 37
4.7. Odd-dimensional spheres 40
References 41

Paul Baum thanks Dartmouth College for the generous hospitality provided to him via the Edward Shapiro fund. Erik van Erp thanks Penn State University for a number of productive and enjoyable visits.

PFB was partially supported by NSF grant DMS-0701184.

EvE was partially supported by NSF grant DMS-1100570.

1. Introduction

1.1. **Context.** By suitably combining analysis and algebraic topology, Michael F. Atiyah and Isodore M. Singer proved their celebrated index theorem for elliptic operators on closed manifolds [2]. In this paper we shall extend the Atiyah-Singer formula to a naturally arising class of *hypoelliptic* (but not elliptic) operators on contact manifolds. The novelty of this paper is that it combines Van Erp's earlier partial results with the Baum-Douglas isomorphism of analytic and geometric K-homology. We find an explicit topological expression for the element in K-homology that corresponds to a hypoelliptic Fredholm operator.

The result is of interest in two directions: (1) it clarifies the index problem itself, resolving some heretofore confusing issues, and (2) it provides interesting new examples of K-cycles that exhibit phenomena not encountered in elliptic theory. The index theorem for elliptic operators has seen numerous generalizations in the context of analytic K-homology and KK-theory, and the theory of hypoelliptic operators can be explored in all of these directions. The identification of the K-cycle corresponding to a hypoelliptic operator (and the resulting formula for the Chern character of the operator) is a key step in this program. For example, it should be possible to derive equivariant and families index formulas for hypoelliptic operators now that we have an expression for the K-cycle. This was not within reach earlier.

A significant advance of index theory from elliptic to hypoelliptic operators was achieved by Alain Connes and Henri Moscovici. In [14] Connes and Moscovici derived an abstract index formula for the Chern character (in cyclic cohomology) of the K-homology element determined by a generalized Fredholm operator (a so-called 'regular spectral triple'). They applied their abstract formula to derive an explicit expression for the Chern character of the transverse fundamental class of a foliated manifold [15]. The Fredholm operator in this spectral triple was not an elliptic but a *hypoelliptic* operator. Hypoelliptic operators were introduced in this context by Michel Hilsum and Georges Skandalis [19], and Connes-Moscivici simplified the construction by working within the *Heisenberg calculus* of [10]. Building on the Connes-Moscivici results, Raphael Ponge proved that the abstract local index formula in cyclic cohomology also applied to hypoelliptic operators in the Heisenberg calculus on *contact manifolds* [22]. Ponge posed (but did not solve) the problem of converting this abstract result into an explicit formula in terms of characteristic classes on the contact manifold. Around the same time, in a context not related to noncommutative geometry, Charles Epstein and Richard Melrose independently investigated a similar index problem for hypoelliptic operators in the Heisenberg calculus on contact manifolds [20]. Their results were reported in [17].

In [24, 25] Van Erp presented an alternative approach to index theory of hypoelliptic operators by adapting the tangent groupoid methods of Connes [16, II.5] to the Heisenberg calculus. The main innovation in [24] was the construction of a K-theory element in $K^0(T^*X)$ for an invertible symbol in the Heisenberg calculus. The symbolic calculus of the Heisenberg algebra is noncommutative, and the principal symbol naturally defines an element in the K-theory of a suitable C^*-algebra related to the calculus. This K-theory group is (via the Connes-Thom isomorphism) naturally isomorphic to $K^0(T^*X)$. Building on this idea, an explicit index formula for scalar (pseudo)differential operators on contact manifolds was derived in [25], and the same general method was applied to derive an index formula for hypoelliptic operators on foliated manifolds in [27].

A curious feature of the index formula in [25] is that it does *not* apply to operators that act on sections of a vector bundle. The method of proof relies on a trick that does not apply in the presence of vector bundles. If P is a hypoelliptic scalar differential operator on a closed contact manifold X then the formula of [25] computes the Fredholm index of P. But if we simply insert the Chern character $\mathrm{ch}(F)$ of a complex vector bundle $F \to X$ in this formula we do not obtain a correct expression for the index of the twisted operator $P \otimes I_F$ acting on sections in F. Put differently, the results in [25] do *not* give a correct expression for the Chern character of the element in K-homology determined by P. For the same reason, the results of [24, 25] do not indicate how to solve the related equivariant anf families index problems. In this sense the result of [25] was partial.

In the current paper we present what we regard as a definitive solution to this index problem on contact manifolds by computing the K-homology class represented by a hypoelliptic Fredholm operator. More precisely, we construct a geometric K-cycle that is equivalent to the analytic K-cycle obtained directly from the operator. From our point of view this fully settles the index problem for the Heisenberg calculus on contact manifolds, and we believe that the result can be generalized to solve the equivariant and families index problems.

1.2. **The problem.** For the purpose of this introduction we discuss a well-known class of second order hypoelliptic operators. In all that follows X will denote a closed smooth contact manifold of dimension $2n+1$. A contact form θ on X is an $\mathbb{R}$-valued 1-form such that $\theta(d\theta)^n$ is a volume form on X. The distribution $H \subset TX$ of codimension 1 that is the kernel of the 1-form θ is called the contact hyperplane bundle. The Reeb vector field T is the unique vector field on X with the properties

$$\theta(T) = 1,\ d\theta(T, \cdot) = 0.$$

The Reeb field T is tranversal to the contact hyperplane bundle H. The 2-form $d\theta$ restricts to a nondegenerate skew-symmetric form in the fibers of H. Thus, the fibers of H are symplectic vector spaces. We choose a complex structure J for H that is compatible with the symplectic form $d\theta$ in each fiber, i.e.,

$$J^2 = -1,\ \ d\theta(Jv, Jw) = d\theta(v,w),\ \ d\theta(Jv, v) \geq 0.$$

Such a choice of J is always possible, because the space of compatible complex structures on a symplectic vector space is contractible. The choice of J also defines a Euclidean structure in the fibers of H, with inner product defined by $\langle v, w\rangle = d\theta(Jv, w)$. In addition, J and θ determine a hermitian inner product,

$$\langle v, w\rangle = d\theta(Jv, w) + i\, d\theta(v, w).$$

A *sublaplacian* for the given structure (X, θ, J) is a second order differential operator on X constructed as follows. Locally, a sublaplacian is of the form

$$\Delta_H = \sum_{j=1}^{2n} -W_j^2.$$

Here $W_1, \ldots, W_{2n}$ is a collection of vector fields, defined in an open subset $U \subseteq X$, such that at each point $x \in U$ the vectors $W_j(x)$ form an orthonormal frame for the Euclidean vector space H_x. A global sublaplacian on X is then constructed via a partition of unity.

To obtain an operator with interesting index theory, we add a first order term tranverse to H and consider second order differential operators of the form

$$P_\gamma = \Delta_H + i\gamma T, \qquad i = \sqrt{-1}$$

where the coefficient γ is a smooth complex-valued function on X. Operators of this type have been studied extensively. See for example [10, Ch. 1] or [17, Thm. 8.1] for the following result.

Proposition 1.2.1. *The operator P_γ is invertible modulo smoothing operators if*

$$\gamma(x) \notin \{\cdots, -n-4, -n-2, -n, n, n+2, n+4, \cdots\}$$

for every point $x \in X$.

The hypoelliptic operator P_γ has essentially the same analytic properties as an elliptic operator. In particular, it gives rise to a cycle in analytic K-homology. The symmetric operator

$$D = \begin{pmatrix} 0 & P_\gamma^* \\ P_\gamma & 0 \end{pmatrix}$$

is essentially self-adjoint, has discrete spectrum, and finite dimensional eigenspaces consisting of smooth functions [24, section 2.6]. The bounded operator $F = D(1 + D^2)^{-1/2}$ defines a graded Fredholm module

$$[P_\gamma] \in KK^0(C(X), \mathbb{C}).$$

Let us briefly verify this. We use the standard convention that $\mathcal{K}$ denotes the separable C^*-algebra of compact operators.

Proposition 1.2.2. *The operator $F = D(1+D^2)^{-1/2}$ is a bounded Fredholm operator on $L^2(X)^{\oplus 2}$ that satisfies the pseudolocality condition $[F, M_f] \in \mathcal{K}(L^2(X)^{\oplus 2})$ for multiplication operators $M_f \in \mathcal{L}(L^2(X)^{\oplus 2}), f \in C(X)$.*

Proof. If P_γ is hypoelliptic then the operator F has order zero in the Heisenberg pseudodifferential calculus. Order zero operators in the Heisenberg calculus are bounded on $L^2(X)^{\oplus 2}$ while operators of negative order are compact (by the Rellich lemma for the Heisenberg calculus). If Ψ_H denotes the norm completion of the algebra of order zero Heisenberg operators, then the principal Heisenberg symbol σ_H gives rise to a short exact sequence of C^*-algebras,

$$0 \to \mathcal{K} \to \Psi_H \overset{\sigma_H}{\to} S_H \to 0.$$

Without knowing the details of the calculus, it suffices to know that the symbol algebra S_H contains $C(X)$ as a central subalgebra. Moreover, for a continuous function $f \in C(X)$ the multiplication operator M_f on $L^2(X)$ is contained in Ψ_H, while its principal symbol is $\sigma_H(M_f) = f$, as usual. It follows that $[\sigma_H(F), \sigma_H(M_f)] = 0$ for any order zero operator F, which is equivalent to $[F, M_f] \in \mathcal{K}$.

□

Our aim is to solve the index problem for hypoelliptic operators in the Heisenberg calculus in the terms formulated by Paul Baum and Ron Douglas in [6].

> *Let X be a closed contact manifold. Given a hypoelliptic (pseudo)differential operator P in the Heisenberg calculus on X representing a cycle in analytic K-homology $[P] \in KK^0(C(X), \mathbb{C})$, explicitly compute an (M, E, φ)-cycle in geometric K-homology $K_0^{top}(X)$ with $\mu(M, E, \varphi) = [P]$.*

Geometric K-cycles were introduced in [5,6] to provide a topological description of K-homology. For a topological space X, a geometric K-cycle is a triple (M, E, φ) consisting of a closed Spin^c manifold M, a complex vector bundle $E \to M$, and a continuous map $\varphi : M \to X$. The collection of (M, E, φ)-cycles, subject to a certain equivalence relation, forms an abelian group under disjoint union. We denote this group by $K_*^{top}(X)$.

There is a natural map

$$\mu : K_j^{top}(X) \to KK^j(C(X), \mathbb{C}), \qquad j = 0, 1$$

from geometric K-homology to analytic K-homology, defined as follows. Let $D_M \otimes I_E$ denote the Dirac operator D_M for the Spin^c manifold M twisted by the vector bundle E. Then $\mu(M, E, \varphi)$ is the push-forward by φ of the analytic K-cycle $[D_M \otimes I_E] \in KK^j(C(M), \mathbb{C})$,

$$\mu(M, E, \varphi) = \varphi_*(D_M \otimes I_E) \in KK^j(C(X), \mathbb{C}).$$

For a finite CW-complex X the map μ is an isomorphism [7].

The point of view of [6] is that index theory, in general, is based on the equivalence between geometric K-homology and analytic K-homology. In [6, section 25] Baum and Douglas ask

> *Let* X *be a finite* CW*-complex. Given* $\xi \in KK^j(C(X), \mathbb{C})$, *explicitly compute the unique* $\tilde{\xi} \in K_j^{top}(X)$ *with* $\mu(\tilde{v}) = v$.

They explain how the solution of several index problems (including, of course, the Atiyah-Singer index formula for elliptic operators) can be understood in this framework. Our aim in this paper is to show that the index problem for the Heisenberg calculus has a very explicit and natural solution when understood in this general context proposed by Baum and Douglas.

If X is a finite CW complex and ξ is an element in $KK^0(C(X), \mathbb{C})$,

$$\xi \in KK^0(C(X), \mathbb{C}),$$

then ξ determines an element $\mathcal{I}(\xi) \in H_{ev}(X; \mathbb{Q})$ with the property: Whenever F is a $\mathbb{C}$ vector bundle on X,

$$\text{Index}\,(F \otimes \xi) = \epsilon_*(\text{ch}(F) \cap \mathcal{I}(\xi)).$$

Here $H_{ev}(X; \mathbb{Q})$ is the direct sum of the even rational homology groups of X,

$$H_{ev}(X; \mathbb{Q}) = H_0(X; \mathbb{Q}) \oplus H_2(X; \mathbb{Q}) \oplus H_4(X; \mathbb{Q}) \oplus \cdots$$

and $\epsilon : X \to \bullet$ is the map of X to a point and $\epsilon_* : H_*(X; \mathbb{Q}) \to H_*(\bullet; \mathbb{Q}) = \mathbb{Q}$ is the resulting map in rational homology. As usual in algebraic topology ch is the Chern character and $\cap$ is cap product. We shall refer to $\mathcal{I}(\xi)$ as the index class of ξ. Equivalently, $\mathcal{I}(\xi)$ is the homology Chern character of ξ. To solve the index problem for ξ *rationally* is to give an explicit formula for $\mathcal{I}(\xi)$. To solve the index problem *integrally* for ξ is to explicitly construct a K-cycle (M, E, φ) on X with

$$\mu(M, E, \varphi) = \xi.$$

If such an (M, E, φ) has been constructed, then

$$\mathcal{I}(\xi) = \varphi_*(\text{ch}(E) \cup \text{Td}(M) \cap [M])$$

where $[M]$ is the fundamental cycle (in $H_*(M; \mathbb{Z})$) of M and $\varphi_* : H_*(M; \mathbb{Q}) \to H_*(X; \mathbb{Q})$ is the map of rational homology induced by $\varphi : M \to X$.

Starting with $\xi \in KK^0(C(X), \mathbb{C})$ the K-cycle (M, E, φ) which integrally solves the index problem for ξ should be constructed so that *mutatis mutandis* the same K-cycle will solve the equivariant index problem (when a compact Lie group is acting) and the index problem for families. We believe that our K-cycle solution for the class of operators considered in this paper is sufficiently natural so that it will also solve the equivariant and families problems.

Now that we have found a geometric K-cycle that solves the problem we finally understand the obstacle that prevented us from finding a solution for the vector bundle problem in our earlier attempts. As we will see, the geometric K-cycle for hypoelliptic operators in the Heisenberg calculus consists of *two* components that cannot be merged, in a natural way, into a single cycle. The K-theoretic methods of [24, 25] conditioned us to look for a *single* expression, but the index formula we find here consists of *two* terms. The Spin^c manifold M in the K-cycle we obtain is familiar in the theory of contact manifolds. It is a compactification of the standard symplectic cone $N^* \smallsetminus 0$ as a submanifold of T^*X with its canonical symplectic structure. Here $N^* \subset T^*X$ denotes the annihilator of H. $N^* \smallsetminus 0$ consists of two copies of $X \times \mathbb{R}$ with opposite symplectic structures. The fact that the final index formula for hypoelliptic operators breaks up into two separate expressions is easily and naturally accomodated by the formalism of geometric

K-homology. Expressing an equivalent solution in the K-theoretic framework of [24, 25] would not be impossible, but the resulting expression would be cumbersome and unnatural.

In summary, K-theory *alone* did not produce a satisfactory answer for this index problem. It was necessary to introduce and use K-homology in order to arrive at a clearly stated solution.

1.3. **Statement of the result.** To make all of this more precise we now discuss the details of the solution of the index problem for the hypoelliptic operators $P_\gamma = \Delta_H + i\gamma T$. The main result of this paper is the solution of the index problem for general hypoelliptic operators in the Heisenberg calculus on contact manifolds. But for the sake of this introduction we restrict our discussion to operators of the form

$$P_\gamma = \Delta_H + i\gamma T.$$

The ingredients of a geometric K-cycle $[(M, E, \varphi)] = \mu^{-1}(P_\gamma)$ are as follows.

The Spin^c manifold M. As mentioned above, a contact manifold has two canonical Spin^c structures. The tangent bundle splits as $TX = H \oplus N$, where N denotes the normal bundle $N = TX/H$. If we choose a global contact 1-form θ, then the normal bundle is trivialized $N = X \times \mathbb{R}$, and it has a Spin^c structure by virtue of this orientation. The orientation of the normal bundle is referred to as the *co-orientation* of the contact structure H. The contact hyperplane bundle H has a Spin^c structure associated to a complex structure J that is compatible, in each fiber, with the symplectic form $d\theta$. Any two choices of J (compatible with θ) result in equivalent Spin^c structures on H. Let us denote by $H^{1,0}$ the contact hyperplane bundle H equipped with its complex structure J, so that $H^{1,0}$ is a complex vector bundle of rank n. The canonical Spin^c structure of X for the choice of contact 1-form θ is, by definition, the product Spin^c structure on the direct sum

$$TX = H \oplus N = H^{1,0} \oplus \underline{\mathbb{R}}.$$

Now consider what happens if we reverse the co-orientation of the contact structure. If we replace θ by $-\theta$ we must also replace the complex structure J with $-J$, because of the compatibility $d\theta(Jv, v) \geq 0$. Thus, the complex structure on H must be conjugated if we reverse the orientation of $N = X \times \mathbb{R}$. The effect on the Spin^c structure of $TX = H \oplus N$ is the combined effect of these two modifications: reversal of orientation of N, and conjugation of the complex structure of $H^{1,0}$. We denote by X^+ the manifold X with its canonical Spin^c structure for the chosen contact form θ, while X^- denotes X with the Spin^c structure for the opposite contact form $-\theta$.

The even dimensional Spin^c manifold M in our (M, E, φ)-cycle is a disjoint union

$$M = X^+ \times S^1 \sqcup X^- \times S^1,$$

with one copy of $X \times S^1$ for each co-orientation of the contact structure. As smooth manifolds $X^+ \times S^1$ and $X^- \times S^1$ are the same, but these two copies of $X \times S^1$ have different Spin^c structures. We shall say that these two Spin^c structures on $X \times S^1$ are *conjugates.*

The fiber of the tangent space $T(X \times S^1) = TX \times TS^1$ is $H_x \times \mathbb{R} \times \mathbb{R}$. We can extend the complex structure J from H_x to $H_x \times \mathbb{R} \times \mathbb{R}$ by identifying $\mathbb{R} \times \mathbb{R} = \mathbb{R} \oplus i\mathbb{R} = \mathbb{C}$. The product Spin^c structure on $X^+ \times S^1$ is then the same as the canonical Spin^c structure corresponding to this complex structure on $T(X \times S^1)$. The product Spin^c structure on $X^- \times S^1$ is the same as the Spin^c structure associated to the *conjugate* complex structure on $T(X \times S^1)$. Therefore one way to think of the manifold M is as an almost complex manifold which is the disjoint union of two copies of $X \times S^1$ with conjugate almost complex structures.

The complex vector bundle E. The vector bundle $E \to M$ consists, of course, of two vector bundles: one on each copy of $X \times S^1$. The vector bundle $E^+ \to X^+ \times S^1$ is constructed by combining symmetric tensors of $H^{1,0}$ with the 'winding' of the coefficient function γ around the positive integers $n+2j, j=0,1,2,\ldots$. The vector bundle $E^- \to X^- \times S^1$ is constructed by combining symmetric tensors of the conjugate bundle $H^{0,1}$ with the winding of γ around the negative integers $-(n+2j), j=0,1,2,\ldots$.

Quite generally, for a topological space W a continuous function $\sigma : W \to \mathbb{C} \smallsetminus \{0\}$ determines a (complex) line bundle $L(\sigma)$ on $W \times S^1$. We set $S^1 = S^1_+ \cup S^1_-$, where S^1_+ and S^1_- are the upper and lower hemispheres of S^1,

$$S^1 = \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}.$$

Then $L(\sigma)$ is obtained by the clutching construction

$$L(\sigma) = (W \times S^1_+) \times \mathbb{C} \cup_\sigma (W \times S^1_-) \times \mathbb{C},$$

where the identification using σ is

$$\begin{aligned} (w,-1,\lambda) &\sim (w,-1,\lambda) \\ (w,1,\lambda) &\sim (w,1,\sigma(w)\lambda). \end{aligned}$$

Equivalently, given $\sigma : W \to \mathbb{C} \smallsetminus \{0\}$ consider the map

$$\tilde{\sigma} : W \times S^1 \to S^1 \times S^1 : (w,\lambda) \mapsto \left(\frac{\sigma(w)}{|\sigma(w)|}, \lambda \right).$$

Then $L(\sigma)$ is the pull-back using $\tilde{\sigma}$ of the Hopf line bundle ξ on $S^1 \times S^1$,

$$L(\sigma) = \tilde{\sigma}^*(\xi).$$

For the integers $k = \pm(n+2j)$ we write

$$L^\gamma_k = L(\gamma - k).$$

Then on $X^+ \times S^1$ we set

$$E^+ = \bigoplus_{j=0}^{N} \left(L^\gamma_{n+2j} \otimes \varphi^* \mathrm{Sym}^j H^{1,0} \right),$$

while on $X^- \times S^1$ we have

$$E^- = \bigoplus_{j=0}^{N} \left(L^\gamma_{-(n+2j)} \otimes \varphi^* \mathrm{Sym}^j H^{0,1} \right),$$

The integer N must be chosen sufficiently large. For large values of j the line bundles $L^\gamma_{\pm(n+2j)}$ are trivial. The resulting contribution to the geometric K-cycle would then be a boundary, and can therefore be omitted. Only finitely many values of j contribute a nontrivial term.

The continuous map φ. The map $\varphi : M \to X$ is projection on the first factor X for each of the two copies of $X \times S^1$.

Theorem 1.3.1. *Let X be a closed manifold with contact form θ, contact hyperplane bundle H and Reeb field T. Suppose a complex structure J is chosen in the fibers of H compatible with the symplectic form $d\theta$. Let Δ_H be a sublaplacian for the structure (X,θ,J).*

For a hypoelliptic Fredholm operator $P_\gamma = \Delta_H + i\gamma T$ we have

$$[P_\gamma] = \mu(X^+ \times S^1,\, E^+,\, \varphi) + \mu(X^- \times S^1,\, E^-,\, \varphi)$$

Remark 1.3.2. Theorem 1.3.1 immediately implies the index formula derived in [25], but is a much stronger and more precise result. It is interesting to compare the two results, and we briefly discuss how to derive the index formula of [25]. As an immediate corollary of Theorem 1.3.1 we obtain

$$\begin{aligned}\operatorname{Index} P_\gamma = &\sum_{j=0}^{N} \int_X \operatorname{ch}(\gamma - (n+2j)) \wedge \operatorname{ch}(\operatorname{Sym}^j H^{1,0}) \wedge \operatorname{Td}(H^{1,0}) \\ &+ (-1)^{n+1} \sum_{j=0}^{N} \int_X \operatorname{ch}(\gamma + (n+2j)) \wedge \operatorname{ch}(\operatorname{Sym}^j H^{0,1}) \wedge \operatorname{Td}(H^{0,1}).\end{aligned}$$

Here X is oriented (in both integrals) by the volume form $\theta(d\theta)^n$. The factor $(-1)^{n+1}$ for the second term arises from the fact that $X^- \times S^1$ has conjugate almost complex structure from $X^+ \times S^1$, and is of complex dimension $n+1$.

The product $\operatorname{ch}(\operatorname{Sym}^j H^{0,1}) \wedge \operatorname{Td}(H^{0,1})$ that appears in the second term is a characteristic class of $H^{0,1}$, and therefore a polynomial in the Chern classes of $H^{0,1}$. For the individual Chern classes we have

$$c_k(H^{0,1}) = (-1)^k c_k(H^{1,0}).$$

Since the odd Chern character $\operatorname{ch}(\gamma + (n+2j))$ of the scalar function $\gamma + (n+2j)$ is a 1-form, the relevant part of $\operatorname{ch}(\operatorname{Sym}^j H^{0,1})\operatorname{Td}(H^{0,1})$ is in dimension $2n$. We can therefore replace $H^{0,1}$ everywhere by $H^{1,0}$ at the expense of a sign $(-1)^n$, obtaining

$$\operatorname{Index} P_\gamma = \sum_{j=0}^{N} \int_X [\operatorname{ch}(\gamma - (n+2j)) - \operatorname{ch}(\gamma + (n+2j))] \wedge \operatorname{ch}(\operatorname{Sym}^j H^{1,0}) \wedge \operatorname{Td}(H^{1,0}).$$

In [25] this was written as

$$\operatorname{Index} P_\gamma = \int_X \operatorname{ch}(a(P)) \wedge \operatorname{Td}(X^+),$$

where $a(P) \in K^1(X)$ is the (odd) K-theory class

$$a(P) = \bigoplus_{j=0}^{N} \left[\operatorname{Sym}^j H^{1,0}, \frac{\gamma - (n+2j)}{\gamma + (n+2j)} \right] \in K^1(X)$$

and $TX^+ = H^{1,0} \oplus \underline{\mathbb{R}}$ has its positively co-oriented Spin^c structure.

Remark 1.3.3. The index formula in the form in which it was presented in [25] leads one to believe that the K-cycle for $[P_\gamma]$ should consist of a single component $(X \times S^1, E, \varphi)$. The above calculation seems to bear this out, because we successfully combined the two components into one. But this is misleading. As soon as we introduce a complex vector bundle $F \to X$ and twist P_γ by F, Theorem 1.3.1 leads to a formula for the Fredholm index of $F \otimes P_\gamma$ that does not easily reduce to a single integral. It is precisely for this reason that the formalism of geometric K-homology appears better suited to formulate the solution to this index problem than topological K-theory. The 'answer' in [25] took the form of a class in $K^1(X)$. Now $K^1(X)$ and $K_0^{top}(X)$ are isomorphic by Poincaré duality. But there are *two* natural Poincaré dualities, one for each of two conjugate Spin^c structures on X, and it is unnatural to try to express the final answer as a single class in $K^1(X)$.

What then is the explanation for the formula of [25] from our current perspective? Effectively, the trick employed in [25] is equivalent, in the current framework, to taking the *conjugate* of every ingredient of the second component of our (M, E, φ) cycle. We can replace the almost complex manifold $X^- \times S^1$ by its conjugate $X^+ \times S^1$, the complex vector bundle $H^{0,1}$ by the conjugate bundle $H^{0,1}$, and the line bundles $L(\gamma + k)$ by their conjugate bundles $L((\gamma + k)^{-1})$.

The resulting K-cycle is a disjoint union of two cycles with Spin^c manifold $X^+ \times S^1$, and they are easily combined into one.

It turns out that this procedure of conjugating the ingredients of an (M, E, φ) cycle constitutes a natural operation in K-homology. In a separate publication we will show that conjugation is a non-trivial automorphism of the K-homology functor that is *not* plus or minus the identity [8]. Therefore, if we conjugate the second term $(X^- \times S^1, E^-, \varphi)$ in our (M, E, φ) cycle for $[P_\gamma]$ we obtain a non-equivalent K-cycle. But now consider what happens if we are only interested in the Fredholm index of the scalar operator P_γ. The Fredholm index corresponds to the map in K-homology $K_0^{top}(X) \to K_0^{top}(\text{point})$ induced by $X \to$ point. As we will show in [8], conjugation in the group $K_0^{top}(\text{point}) = \mathbb{Z}$ is just the identity map. Therefore, while conjugation of the second term in our K-cycle alters its equivalence class in $K_0^{top}(X)$, this operation does not affect its image in $K_0^{top}(\text{point})$. The discovery and clarification of the conjugation operation in K-homology is an interesting spin-off of our current work. We refer to [8] for a detailed discussion of this operation.

1.4. **Outline of the proof.** So far we have only considered second order differential operators $P_\gamma = \Delta_H + i\gamma T$. But our results apply to arbitrary hypoelliptic pseudodifferential operators in the Heisenberg calculus. In this section we outline the general results and the main structure of the proof. The following diagram represents the overall structure of the argument.

$$\begin{tikzcd}
 & K_0(C^*(T_H X)) \arrow[ddl, bend right, "b"'] \arrow[ddr, bend left, "\mathrm{Op_H}"] & \\
 & K^0(T^*X) \arrow[u, "\Psi"] \arrow[dl, "c"] \arrow[dr, "\mathrm{Op_e}"'] & \\
K_0^{top}(X) \arrow[rr, "\mu"'] & & KK^0(C(X), \mathbb{C})
\end{tikzcd}$$

All the arrows in this diagram are isomorphisms. Theorem 1.3.1 will follow as soon as we have shown that the outer triangle commutes. We prove that the outer triangle commutes by showing that the *other* three triangles in the diagram commute. Commutativity of each of these triangles constitutes an independent result, i.e., the three commutativities are logically independent.

The Atiyah-Singer index theorem. The bottom triangle in the diagram,

$$\begin{tikzcd}
 & K^0(T^*X) \arrow[dl, "c"'] \arrow[dr, "\mathrm{Op_e}"] & \\
K_0^{top}(X) \arrow[rr, "\mu"'] & & KK^0(C(X), \mathbb{C})
\end{tikzcd}$$

first appeared in [6, section 23], and presents the index theorem of Atiyah and Singer as a statement in K-homology. The map

$$c : K^0(T^*X) \to K_0^{top}(X)$$

is the 'clutching construction' of [6, section 22], which produces an (M, E, φ)-cycle from a (compactly supported) K-theory class $(\sigma, E^0, E^1) \in K^0(T^*X)$. One can think of this map as implementing Poincaré duality between $K^0(T^*X)$ and $K_0^{top}(X)$.

The map

$$\mathrm{Op}_e : K^0(T^*X) \to KK^0(C(X), \mathbb{C})$$

is the 'choose an elliptic operator' map: For a compactly supported K-theory class (σ, E^0, E^1) one constructs an order zero pseudodifferential elliptic operator $P : C^\infty(X, E^0) \to C^\infty(X, E^1)$ whose symbol (at the level of K-theory) is (σ, E^0, E^1). Commutativity of the triangle amounts to the statement that the (M, E, φ)-cycle obtained by clutching the symbol of an elliptic operator P is the topological equivalent of the analytic K-cycle $[P]$. This is the Atiyah-Singer index theorem for elliptic operators formulated in the context of geometric K-homology. This was essentially stated and proved in [1]. The proof implicit in [1] uses the Atiyah-Singer theorem for families of elliptic operators. Elsewhere [9] we shall give a proof of the commutativity of this diagram which does not use the Atiyah-Singer index theorem. Thus the situation shall be reversed. In [9] Atiyah-Singer will be a corollary of commutativity of the triangle.

The index theorem for the Heisenberg calculus. The triangle

$$\begin{tikzcd}
 & K_0(C^*(T_H X)) \arrow[dl, "b"'] \arrow[dr, "\mathrm{Op}_{\mathrm{H}}"] & \\
K_0^{top}(X) \arrow[rr, "\mu"'] & & KK^0(C(X), \mathbb{C})
\end{tikzcd}$$

is the analog for the index problem of hypoelliptic operators on contact manifolds of the triangle of [6, section 23].

Elements in the group $K^0(T^*X)$ represent symbols of elliptic operators. For the Heisenberg calculus the group $K^0(T^*X)$ needs to be replaced by $K_0(C^*(T_H X))$, the K-theory of a noncommutative C^*-algebra $C^*(T_H X)$. The notation $T_H X$ refers to the tangent bundle TX equipped with some extra structure related to the contact form. The symplectic form on H_x gives rise to the structure of a Heisenberg group in each tangent fiber $T_x X = H_x \times \mathbb{R}$, with product

$$(v, t)(v', t') = (v + v', t + t' - \frac{1}{2} d\theta(v, v')), \quad v, v' \in H_x,\ t, t' \in \mathbb{R}.$$

Thus, $T_H X$ is a fiber bundle of nilpotent Lie groups instead of a bundle of vector spaces. We will think of $T_H X$ as a smooth groupoid with base X. The C^*-algebra $C^*(T_H X)$ is the norm completion of the convolution algebra $C_c^\infty(T_H X)$ of functions on the groupoid $T_H X$.

The insight that $K_0(C^*(T_H X))$ is the natural recipient for the principal symbol of a Fredholm operator in the Heisenberg calculus is a central result in [24]. We will show here that there is a well-defined map

$$\mathrm{Op}_H : K_0(C^*(T_H X)) \to KK^0(C(X), \mathbb{C})$$

which is the 'choose a hypoelliptic operator' map for the Heisenberg calculus: To a class $v \in K_0(C^*(T_H X))$ assign an order zero hypoelliptic operator in the Heisenberg calculus whose principal Heisenberg symbol is equivalent in K-theory to v.

Explicitly identifying the map

$$b : K_0(C^*(T_H X)) \to K_0^{top}(X)$$

that makes the triangle commute is the crux of our index problem. Given the fact that $C^*(T_H X)$ is noncommutative, the definition of b is necessarily more complicated than in the classical elliptic case (see Definition 2.7.1). Commutativity of the triangle is the content of Theorem 3.5.1, which is the main result of this paper. The crucial point is, of course, that the Poincaré duality map b can be effectively computed, which we show in the remainder of section 4.

The maps to geometric K-homology. We prove our index theorem by showing that the two remaining triangles in our overall diagram commute. Because $T_H X$ is a bundle of nilpotent groups, there is a natural isomorphism

$$\Psi : K^0(T^*X) \to K_0(C^*(T_HX)).$$

The isomorphism Ψ ultimately derives from the Connes-Thom isomorphism [21]. The remaining commutative triangles concern the compatibility of Ψ with the maps to the two K-homology groups. One of the two remaining triangles is

$$\begin{tikzcd}
K_0(C^*(T_HX)) \arrow[dr, bend left, "b"] & \\
K^0(T^*X) \arrow[u, "\Psi"] \arrow[r, "c"] & K_0^{top}(X)
\end{tikzcd}$$

Commutativity of this triangle is the content of Theorem 2.7.5. The map b is the analog in the context of the Heisenberg calculus of the 'clutching construction' c of [6], which is a version of Poincaré duality. Indeed, our construction of the map b involves Poincaré duality in an essential way. In fact, it involves Poincaré duality for *both* of the natural Spin^c structures associated to the contact structure. Thus, Poincaré duality (and the fact that there are *two* such dualities for every contact manifold) is at the heart of our solution of the index problem on contact manifolds.

Proving that the triangle commutes involves effectively computing an inverse for the Connes-Thom isomorphism Ψ. This is one of the most difficult steps in the proof, and it was not achieved in either [24] or [25]. We succeed here by combining a careful analysis of the Type I structure of $C^*(T_HX)$ with a strong deformation of this noncommutative algebra to $C_0(T^*X)$.

The maps to analytic K-homology. The link between the two 'choose an operator' maps for the classical and Heisenberg calculi is established in the triangle:

$$\begin{tikzcd}
K_0(C^*(T_HX)) \arrow[dr, bend left, "\mathrm{Op_H}"] & \\
K^0(T^*X) \arrow[u, "\Psi"] \arrow[r, "\mathrm{Op_e}"'] & KK^0(C(X),\mathbb{C})
\end{tikzcd}$$

Using an adaptation of the tangent groupoid formalism of Alain Connes, we showed in [24, section 3.7] that the isomorphism Ψ commutes with the analytic index maps,

$$\mathrm{Index}\,\mathrm{Op}_e(\sigma) = \mathrm{Index}\,\mathrm{Op_H}(\Psi(\sigma)).$$

In other words, we established commutativity of

$$\begin{tikzcd}
K_0(C^*(T_HX)) \arrow[dr, bend left, "\mathrm{IndexOp_H}"] & \\
K^0(T^*X) \arrow[u, "\Psi"'] \arrow[r, "\mathrm{IndexOp_e}"'] & \mathbb{Z}
\end{tikzcd}$$

where the 'choose an operator maps' were composed with the Fredholm index

$$\mathrm{Index} : KK^0(C(X),\mathbb{C}) \to \mathbb{Z}.$$

Here we modify the tangent groupoid argument from [24] and derive the stronger result that

$$[\mathrm{Op_e}(\sigma)] = [\mathrm{Op_H}(\Psi(\sigma))] \in KK(C(X),\mathbb{C}).$$

This is the content of Theorem 3.4.1.

1.5. **Summary of the paper.** Section 2 discusses the ingredients and commutativity of the triangle

$$\begin{tikzcd}
K_0(C^*(T_H X)) \arrow[rd, bend left, "b"] & \\
K^0(T^*X) \arrow[u, "\Psi"] \arrow[r, "c"] & K_0^{top}(X)
\end{tikzcd}$$

A major portion of this section consists of a detailed analysis of the structure of the C^*-algebra $C^*(T_H X)$. This analysis culminates in Proposition 2.6.1, which is the basis for our Definition 2.7.1 of the 'Poincaré duality' map b from K-theory $K_0(C^*(T_H X))$ to K-homology $K_0^{top}(X)$. Theorem 2.7.5 is the conclusion of this section, establishing commutativity of the triangle above.

Section 3 deals with the analogous triangle for analytic K-homology,

$$\begin{tikzcd}
K_0(C^*(T_H X)) \arrow[rd, bend left, "\mathrm{Op_H}"] & \\
K^0(T^*X) \arrow[u, "\Psi"] \arrow[r, "\mathrm{Op_e}"'] & KK^0(C(X),\mathbb{C})
\end{tikzcd}$$

resulting in Theorem 3.4.1 which establishes commutativity of the triangle. Here we also show that the 'choose a hypoelliptic operator' map $\mathrm{Op_H}$ is well-defined. Combining the results of sections 2 and 3 yields what we consider to be our main result: Theorem 3.5.1 which asserts commutativity of the triangle

$$\begin{tikzcd}
& K_0(C^*(T_H X)) \arrow[ld, "b"'] \arrow[rd, "\mathrm{Op_H}"] & \\
K_0^{top}(X) \arrow[rr, "\mu"'] & & KK^0(C(X),\mathbb{C})
\end{tikzcd}$$

In section 4 we show how to apply Theorem 3.5.1 to compute an explicit K-cycle $b(\sigma_H(P)) = (M, E, \varphi)$ for a given hypoelliptic operator P in the Heisenberg calculus. In section 4.3 we carry out this computation for the special case of the operators $P_\gamma = \Delta_H + i\gamma T$, proving Theorem 1.3.1. In section 4.4 we indicate how the Boutet de Monvel index formula for Toeplitz operators is a corollary of Theorem 3.5.1. In section 4.6 we answer a question posed by R. Ponge. We conclude this paper with the example of odd-dimensional spheres, where our index formula has an especially simple form.

Acknowledgment. We thank A. Connes, R. G. Douglas, E. Getzler, A. Gorokhovsky, N. Higson, R. Nest, and R. Willett for enlightening comments and discussions.

2. The Poincaré duality map b

This section is an exercise in noncommutative topology. We will not yet be concerned with hypoelliptic operators. Instead we focus here on a thorough analysis of the structure of the groupoid C^*-algebra $C^*(T_H X)$. This then allows us to construct an inverse for the Connes-Thom isomorphism

$$\Psi\colon K^0(T^*X) \to K_0(C^*(T_H X)).$$

In Definition 2.7.1 we define the Poincaré duality map

$$b\colon K_0(C^*(T_H X)) \to K_0^{top}(X).$$

The main result of this section is Theorem 2.7.5, which asserts that there is commutativity in the triangle

$$\begin{tikzcd}
K^0(T^*X) \arrow[r, "\Psi"] \arrow[d, "c"] & K_0(C^*(T_HX)) \arrow[dl, "b"] \\
K_0^{top}(X) &
\end{tikzcd}$$

The Type I C^*-algebra $C^*(T_HX)$ decomposes into stably commutative factors. In K-theory each factor in this decomposition can be treated as a topological space. This is our first handle on the K-theory of $C^*(T_HX)$.

Deformations, if appropriately axiomatized, induce maps in K-theory. Alain Connes's *tangent groupoid* proof of the Atiyah-Singer theorem beautifully encodes the role of C^*-algebra deformations in index theory [16, II.5]. In [24] we adapted the tangent groupoid proof to the hypoelliptic index problem. Our strategy for computing the topological equivalent for an element in $K_0(C^*(T_HX))$ is to realize the decomposition of $C^*(T_HX)$ into stably commutative factors as the quantization of a purely commutative extension.

2.1. **The C^*-algebra of the Heisenberg group.** The C^*-algebra $C^*(T_HX)$ is a locally trivial bundle of C^*-algebras over X, whose fiber is the group C^*-algebra $C^*(G)$ of the Heisenberg group. We start by recalling some well-known facts about the structure of this algebra.

Let $V = \mathbb{R}^{2n}$ be the standard symplectic vector space with symplectic form ω. Multiplication in the Heisenberg group $G = V \times \mathbb{R}$ is defined by

$$(v,t)\cdot(v',t') = (v+v', t+t' + \frac{1}{2}\omega(v,v')).$$

The factor $\frac{1}{2}$ is chosen so that for two vectors $v, w \in V$ the Lie bracket is given by the symplectic form $[v,w] = \omega(v,w)$.

Fourier transform in the $\mathbb{R}$ variable for functions $f \in C_c^\infty(G)$,

$$\tilde{f}(v,s) = \int_{\mathbb{R}} e^{-its} f(v,t)dt$$

completes to a $*$-isomorphism,

$$C^*(G) \cong C^*(V \times \mathbb{R}^*, s\omega).$$

Here $V \times \mathbb{R}^*$ denotes the vector bundle over $\mathbb{R}^*$ with fiber V. We may think of it as a smooth groupoid with object space $\mathbb{R}^*$. The expression $s\omega$ refers to the groupoid cocycle

$$c\colon V \times V \times \mathbb{R}^* \to U(1)\ ,\ c((v,s),(w,s)) = \exp\left(is\omega(v,w)\right).$$

The C^*-algebra $C^*(V\times\mathbb{R}^*, s\omega)$ is the completion of the groupoid convolution algebra $C_c^\infty(V\times\mathbb{R}^*)$ twisted by the cocycle c,

$$(\tilde{f} *_c \tilde{g})(v,s) = \int_V e^{is\omega(v,w)} \tilde{f}(v-w,s)\tilde{g}(w,s)\, dw.$$

Elements in the C^*-algebra $C^*(G)$ can be identified quite naturally with sections in a continuous field over $\mathbb{R}^*$. The fiber at $s \in \mathbb{R}^*$ is the twisted convolution C^*-algebra $C^*(V, s\omega)$ (now with fixed value of the parameter s appearing in the cocycle). At $s = 0$ the fiber is commutative, $C^*(V) \cong C_0(V^*)$. If $s \neq 0$ the field is trivial, and each fiber is isomorphic to the algebra of compact operators, $C^*(V, s\omega) \cong \mathcal{K}$. Restriction to $s = 0$ gives the decomposition

$$0 \to C^*(V \times \mathbb{R}^\times, s\omega) \to C^*(G) \to C_0(V^*) \to 0,$$

A simple rescaling of the parameter $s \in \mathbb{R}^\times = \mathbb{R}^* \setminus \{0\}$ gives an isomorphism

$$C^*(V \times \mathbb{R}^\times, s\omega) \cong C_0(-\infty, 0) \otimes C^*(V, -\omega) \oplus C_0(0, \infty) \otimes C^*(V, \omega).$$

We adopt the convention that we identify

$$X \times \mathbb{R}^\times = X \times (-\infty, 0) \sqcup X \times (0, \infty) \approx X \times \mathbb{R} \sqcup X \times \mathbb{R}$$

by the map $s \mapsto \log|s|$ for $s \in \mathbb{R}^\times$. In other words, both components of $X \times \mathbb{R}^\times$ are oriented from 0 to $\pm\infty$. In then have an isomorphism

$$C^*(V \times \mathbb{R}^\times, s\omega) \cong C_0(\mathbb{R}) \otimes C^*(V, -\omega) \oplus C_0(\mathbb{R}) \otimes C^*(V, \omega).$$

While both C^*-algebras $C^*(V, \pm\omega)$ are isomorphic to $\mathcal{K}$ we will not identify these two algebras, for reasons that will soon become apparent.

2.2. $U(n)$ **symmetry.** With a choice of contact 1-form θ and compatible complex structure J in the fibers of H we have a reduction of the structure group of TX to $U(n)$. It is useful to consider the $U(n)$ symmetry of the group C^*-algebra $C^*(G)$.

With the standard identification $V = \mathbb{R}^{2n} = \mathbb{C}^n$ the canonical action of the unitary group $U(n)$ on $V = \mathbb{C}^n$ induces automorphisms of the Heisenberg group $G = V \times \mathbb{R}$, and therefore of the C^*-algebra $C^*(G)$. There is also the obvious action of $U(n)$ on the C^*-algebra $C^*(V \times \mathbb{R}^*, s\omega)$. The $U(n)$ action on functions on $G = V \times \mathbb{R}$ commutes with the Fourier transform in the $\mathbb{R}$ variable. Therefore the isomorphism

$$C^*(G) \cong C^*(V \times \mathbb{R}^*, s\omega)$$

is $U(n)$ equivariant. Likewise, the isomorphism

$$C^*(V \times \mathbb{R}^\times, s\omega) \cong C(\mathbb{R}) \otimes C^*(V, \omega) \oplus C(\mathbb{R}) \otimes C^*(V, -\omega)$$

is $U(n)$ equivariant.

Now choose a representation

$$C^*(V, \omega) \cong \mathcal{K}(\mathcal{H}).$$

The $U(n)$ action on $C^*(V, \omega)$ induces an action on $\mathcal{K}(\mathcal{H})$. The automorphism group of $\mathcal{K}(\mathcal{H})$ is the projective unitary group $U(\mathcal{H})/U(1)$ of the separable Hilbert space $\mathcal{H}$. As we will see below, the induced action $U(n) \to \mathrm{Aut}(\mathcal{K}(\mathcal{H}))$ lifts to a unitary representation $U(n) \to U(\mathcal{H})$. This is of crucial importance for our analysis, because it will imply, later on, the vanishing of the Dixmier-Douady invariant of a specific $\mathcal{K}(\mathcal{H})$-bundle over X, providing us with an important Morita equivalence.

For the opposite algebra $C^*(V, -\omega)$ we have the isomorphism

$$C^*(V, -\omega) \cong \mathcal{K}(\overline{\mathcal{H}}),$$

where $\overline{\mathcal{H}}$ denotes the Hilbert space that is conjugate (or dual) to $\mathcal{H}$, and that carries the dual representation of $U(n)$. This isomorphism is equivariant with respect to the $U(n)$ actions, and the Hilbert space $\overline{\mathcal{H}}$ is *not* ismorphic to $\mathcal{H}$ *as a representation space* of $U(n)$.

In summary, we obtain the $U(n)$ equivariant isomorphism

$$C^*(V \times \mathbb{R}^\times, s\omega) \cong C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}) \oplus C_0(\mathbb{R}) \otimes \mathcal{K}(\overline{\mathcal{H}}).$$

It will be useful to have a concrete model for the Hilbert space $\mathcal{H}$ with its representations of the Heisenberg group G and the unitary group $U(n) \subset \mathrm{Aut}(G)$.

2.3. **Bargmann-Fok space.** The representation space of the Heisenberg group G that most clearly exhibits the $U(n)$ action is the Bargmann-Fok space. As a $U(n)$ representation, the Bargmann-Fok space is a completion of the space of symmetric tensors of $\mathbb{C}^n$,

$$\mathcal{H}^{BF} \supset \operatorname{Sym}\mathbb{C}^n = \bigoplus_{j=0}^{\infty} \operatorname{Sym}^j \mathbb{C}^n$$

with its standard $U(n)$ action. If we identify symmetric tensors with complex polynomials on $\mathbb{C}^n$ then the Bargmann-Fok space is the Hilbert space of entire functions on $\mathbb{C}^n$ with inner product

$$\langle f, g\rangle = \int f(z)\overline{g(z)}\, e^{-|z|^2} dz.$$

If $Z_1, \ldots, Z_n$ is an orthonormal basis for $\mathbb{C}^n$ then the monomials $\Pi_{i=1}^n Z_i^{m_i}/m_i!$ form an orthonormal basis for $\mathcal{H}^{BF}$. In particular, the summands $\operatorname{Sym}^j \mathbb{C}^n$ are mutually orthogonal.

Let us briefly review how the Heisenberg group G acts on this space. To describe the representation it is convenient to pass to the complexified Lie group $G_{\mathbb{C}}$. In order not to get confused, let us denote by $J\colon V \to V$ the standard complex structure on $V = \mathbb{R}^{2n}$ and *not* identify V with $\mathbb{C}^n$ here. The complexified space $V \otimes \mathbb{C}$ splits into the $\pm\sqrt{-1}$ eigenspaces for J as $V \otimes \mathbb{C} = V^{1,0} \oplus V^{0,1}$. Correspondingly, the complexification of the Lie algebra $\mathfrak{g}$ of G splits as a direct sum

$$\mathfrak{g} \otimes \mathbb{C} = V^{1,0} \oplus V^{0,1} \oplus \mathbb{C}.$$

Recall that $\omega(Ju, Jv) = \omega(u, v)$ and $\omega(Jv, v) > 0$ if $v \neq 0$. If we extend the bilinear form ω to a complex bilinear form on $V \otimes \mathbb{C}$ then the expression

$$\langle z, w\rangle = i\omega(z, \bar{w})$$

defines a hermitian inner product on $V^{1,0}$. Let $\{Z_1, \ldots, Z_n\}$ be a basis for $V^{1,0}$ that is orthonormal with respect to this hermitian product. For the complexified Lie algebra $A, B \in \mathfrak{g} \otimes \mathbb{C}$ we have $[A, B] = \omega(A, B) = -i\langle A, \bar{B}\rangle$. Therefore

$$[Z_j, \bar{Z}_k] = \frac{1}{i}\delta_{jk}.$$

Since $V^{1,0}$ is perpendicular to $V^{0,1}$ we also obtain:

$$[Z_j, Z_k] = [\bar{Z}_j, \bar{Z}_k] = 0.$$

In this picture the Bargmann-Fok space contains the symmetric powers of $V^{1,0}$, and can be identified with a space of entire functions on $V^{0,1}$. The representation π of the complexified Lie algebra $\mathfrak{g} \otimes \mathbb{C}$ on $\mathcal{H}^{BF}$ is given by

$$\pi(Z_j) = iz_j \ , \ \pi(\bar{Z}_j) = i\frac{\partial}{\partial z_j} \ , \ \pi(1) = i.$$

The multiplication operator z_j and complex derivative $\partial/\partial z_j$ are unbounded linear operators on $\mathcal{H}^{BF}$. The subspace of polynomials $\operatorname{Sym} V^{1,0}$ is an invariant subspace for both operators, and hence also for the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ of the Lie algebra (they are smooth vectors of the representation). The operators z_j and $\partial/\partial z_j$ are adjoints, and so $\pi(W)^* = -\pi(\bar{W})$ for all vectors $W \in \mathfrak{g} \otimes \mathbb{C}$. This implies that $\pi(v)$ is skew adjoint for vectors in the (real) Lie algebra $v \in \mathfrak{g}$, and the corresponding representation for the group G is unitary.

In summary, we have $U(n)$ equivariant $*$-homomorphisms

$$\begin{tikzcd}
C^*(G) \arrow[d, "f\mapsto\tilde{f}(-{,}1)"'] \arrow[dr, "\pi"] & \\
C^*(V,\omega) \arrow[r, "\cong"] & \mathcal{K}(\mathcal{H}^{BF}).
\end{tikzcd}$$

The horizontal arrow is the Bargmann integral transform, which assigns to a function on V (or its Fourier transform on V^*) an operator on $\mathcal{H}^{BF}$ [4]. For the opposite algebra $C^*(V,-\omega)$ we obtain a representation on the conjugate space $\overline{\mathcal{H}}^{BF}$, which is a completion of the space of symmetric tensors of $V^{0,1} = \overline{\mathbb{C}}^n$ carrying the dual (or conjugate) representation of $U(n)$. We will adopt the notation

$$\mathcal{H}_+^{BF} = \mathcal{H}^{BF},\ \mathcal{H}_-^{BF} = \overline{\mathcal{H}}^{BF}.$$


$\hat{G}$
$\mathcal{K}(\mathcal{H}^{BF}) \cong C^*(V,\omega)$
$C_0(V^*) \cong C^*(V)$
$\mathcal{K}(\overline{\mathcal{H}}^{BF}) \cong C^*(V,-\omega)$

FIGURE 1. The short exact sequence $0 \to C_0(\mathbb{R}^\times, \mathcal{K}) \to C^*(G) \to C_0(V^*) \to 0$

2.4. **Quantization.** Recall that the group C^*-algebra $C^*(G)$ is a quantization of the commutative algebra $C_0(\mathfrak{g}^*) = C_0(V^* \times \mathbb{R}^*)$. The procedure is quite simple. Consider the twisted convolution C^*-algebra

$$A = C^*(V \times \mathbb{R}^* \times [0,1], s\varepsilon\omega),$$

with cocycle

$$c\colon V \times V \times \mathbb{R}^* \times [0,1] \to U(1)\ ,\ c((v,s,\varepsilon),(w,s,\varepsilon)) = \exp(is\varepsilon\omega(v,w)).$$

We can think of elements in this C^*-algebra A as sections in a continuous field $\{A_\varepsilon\}$ with parameter $\varepsilon \in [0,1]$. For $\varepsilon \in (0,1]$ the field is trivial, with $A_\varepsilon \cong A_1 = C^*(G)$. At $\varepsilon = 0$ we have the commutative algebra

$$A_0 = C^*(V \times \mathbb{R}^*) \cong C_0(V^* \times \mathbb{R}^*).$$

We thus have a (strong deformation) quantization from A_0 to A_1. As is well-known, this quantization induces an invertible E-theory element in $E(A_0, A_1)$ that, in turn, corresponds to a KK-equivalence.

Now consider the decomposition

$$0 \to C^*(V \times \mathbb{R}^\times, s\omega) \to C^*(G) \to C^*(V) \to 0.$$

When restricted to the ideal

$$C^*(V \times \mathbb{R}^\times, s\omega) \cong C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}_-^{BF}) \oplus C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}_+^{BF})$$

the KK-equivalence $C^*(G) \sim C_0(\mathfrak{g}^*)$ amounts to the Morita equivalence

$$C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}_-^{BF}) \oplus C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}_+^{BF})\ \sim\ C_0(\mathbb{R}) \otimes \mathbb{C} \oplus C_0(\mathbb{R}) \otimes \mathbb{C}$$

composed with the Bott isomorphisms $\beta_- \oplus \beta_+$,

$$\cdots \sim C_0(\mathbb{R}) \otimes C_0(V^*) \oplus C_0(\mathbb{R}) \otimes C_0(V^*) \cong C_0(V^* \times \mathbb{R}^\times).$$

Here β_+ denotes the canonical Bott generator for the complex vector space $V^{1,0} = \mathbb{C}^n$, while β_- denotes the Bott generator for the conjugate space $V^{0,1}$,

$$\beta_+ \in KK(\mathbb{C}, C_0(V^{1,0})), \;\; \beta_- \in KK(\mathbb{C}, C_0(V^{0,1})).$$

We obtain a KK-equivalence of short exact sequences:

$$\begin{tikzcd}
0 \arrow[r] & C^*(V \times \mathbb{R}^\times, s\omega) \arrow[r] \arrow[d, dotted, "KK^{U(n)}"', "\cong"] & C^*(G) \arrow[r] \arrow[d, dotted, "KK^{U(n)}"', "\cong"] & C^*(V) \arrow[r] \arrow[d, "\cong"] & 0 \\
0 \arrow[r] & C_0(V^* \times \mathbb{R}^\times) \arrow[r] & C_0(V^* \times \mathbb{R}^*) \arrow[r] & C_0(V^*) \arrow[r] & 0
\end{tikzcd}$$

The close connection between Bott periodicity and the KK-equivalence $C^*(V \times \mathbb{R}^\times, s\omega) \sim C_0(V^* \times \mathbb{R}^\times)$ gives us a grip on the KK-equivalence $C^*(G) \sim C_0(V^* \times \mathbb{R}^*)$, which is the Connes-Thom isomorhism that we are trying to understand. Observe that all $*$-homomorphisms and KK-equivalences are $U(n)$-equivariant. The diagram can therefore be interpreted as a commutative diagram in the category $KK^{U(n)}$.

2.5. **Associated bundles.** Our entire analysis easily carries over to associated bundles (of groups, Hilbert spaces, C^*-algebras) over a contact manifold X. As above, the contact 1-form θ and complex structure J in the fibers of H induce a reduction of the structure group of the tangent bundle TX to $U(n)$. Let P_U denote the principal $U(n)$ bundle of orthonormal frames in H with respect to its hermitian structure associated to θ, J.

The standard action α of $U(n)$ on $\mathbb{C}^n$ gives the associated complex vector bundle

$$H^{1,0} = P_U \times_\alpha \mathbb{C}^n.$$

This is just the complex vector bundle with underlying real vector bundle H and complex structure J. The tangent space TX is then identified with

$$TX = P_U \times_{\alpha \oplus 1} \mathbb{C}^n \oplus \mathbb{R} = H^{1,0} \oplus \underline{\mathbb{R}}.$$

This identification exhibits the stably almost complex (and hence Spin^c) structure of TX. The bundle of Heisenberg groups $T_H X$ identifies with

$$T_H X = P_U \times_{\alpha \times 1} G.$$

Here $\alpha \times 1$ denotes the $U(n)$ action on $G = \mathbb{C}^n \times \mathbb{R}$. The induced action on $C^*(G)$ gives the convolution C^*-algebra $C^*(T_H X)$ of the groupoid $T_H X$,

$$C^*(T_H X) = P_U \times_{\alpha \times 1} C^*(G).$$

Denote by ρ the action of $U(n)$ on the Bargmann-Fok space $\mathcal{H}^{BF} = \mathcal{H}^{BF}_+$, the completion of the symmetric tensors $\mathrm{Sym}\,\mathbb{C}^n$ described above. We can form the associated bundle of Hilbert spaces

$$V^{BF}_+ = P_U \times_\rho \mathcal{H}^{BF}_+.$$

Continuous sections in V^{BF}_+ form a Hilbert module over $C(X)$, and we have the dense subspace

$$\bigoplus_{j=0}^{\infty} \mathrm{Sym}^j H^{1,0} \subset V^{BF}_+.$$

The conjugate representation to ρ gives rise to the dual module V_-^{BF} with fiber $\mathcal{H}_-^{BF}$ and

$$\bigoplus_{j=0}^{\infty} \mathrm{Sym}^j H^{0,1} \subset V_-^{BF}.$$

If π denotes the Bargmann-Fok representation of G on $\mathcal{H}^{BF}$, then for fixed $u \in U(n)$ the operator $\rho(u) \in U(\mathcal{H}^{BF})$ is an intertwiner of π and $\pi \circ (\alpha \times 1)(u)$,

$$\pi((\alpha \times 1)(u).g) = \rho(u)\pi(g)\rho(u)^{-1}.$$

This compatibility of $\alpha \times 1$, ρ and π (respectively: $U(n)$ acting on G, $U(n)$ acting on $\mathcal{H}^{BF}$, and G acting on $\mathcal{H}^{BF}$) implies that π is well-defined as a representation of a fiber $C^*(G_x)$ of the bundle $C^*(T_H X)$ on the fiber V_x^{BF} of the Hilbert module V_+^{BF}. The representation π thus gives rise to a $\star$-homomorphism of $C(X)$-algebras,

$$\pi : C^*(T_H X) \to \mathcal{K}(V_+^{BF}).$$

Here $\mathcal{K}$ refers to 'compact operators' in the sense of Hilbert modules. Our analysis of the structure of $C^*(G)$ carries over to $C^*(T_H X)$. We obtain a short exact sequence

$$0 \to I_H \to C^*(T_H X) \to C_0(H^*) \to 0,$$

and the ideal I_H can be identified with

$$\begin{aligned} I_H &\cong P_U \times_{1\otimes Ad(\overline{\rho})} [C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}_-^{BF})] \;\oplus\; P_U \times_{1\otimes Ad(\rho)} [C_0(\mathbb{R}) \otimes \mathcal{K}(\mathcal{H}_+^{BF})] \\ &\cong C_0(\mathbb{R}) \otimes \mathcal{K}(V_-^{BF}) \oplus C_0(\mathbb{R}) \otimes \mathcal{K}(V_+^{BF}). \end{aligned}$$

We see that I_H is Morita equivalent to the commutative algebra $C_0(X \times \mathbb{R}^\times)$, and we have an explicit imprimitivity bimodule, namely the disjoint union of the pull-back of V_+^{BF} to $X \times (0, \infty)$ and the pull-back of V_-^{BF} to $X \times (-\infty, 0)$.

2.6. **Inverting the Connes-Thom isomorphism.** In [24] we discussed the crucial role of the isomorphism

$$\Psi : K^0(T^*X) \to K_0(C^*(T_H X))$$

for our index problem. The isomorphism Ψ is essentially the Connes-Thom isomorphism in each fiber, which agrees with the KK-equivalence induced by the quantization from $C_0(\mathfrak{g}^*)$ to $C^*(G)$. We now obtain a better grip on this isomorphism by analyzing how the quantization behaves when we decompose $C^*(T_H X)$. Our analysis of $C^*(G)$ carries over immediately to $C^*(T_H X)$ because of the $U(n)$ equivariance of all the relevant constructions.

Quite generally, a principal G bundle P for compact group G and compact base $X = P/G$ induces a functor from the category KK^G (equivariant KK-theory) to the category KK^X (RKK-theory for $C(X)$-algebras) in the obvious way. The functor assigns to a G-C^*-algebra A the $C(X)$-C^*-algebra of G-equivariant continuous functions

$$P(A) = \{f : P \to A \mid f \text{ continuous},\ f(pg) = g^{-1}f(p)\}.$$

Similarly for the morphisms: to a G Hilbert module $\mathcal{E}$ over A the functor assigns the $C(X)$ Hilbert module of G-equivariant continuous functions $P \to \mathcal{E}$, etc.

In this way the $U(n)$-equivariant KK-equivalence of short exact sequences for $C^*(G)$ that we derived above implies KK-equivalence of the associated sequences for bundles over X. We obtain

$$\begin{tikzcd}
0 \arrow[r] & I_H \arrow[r] \arrow[d, dotted, "KK^X"', "\cong"] & C^*(T_HX) \arrow[r] \arrow[d, dotted, "KK^X"', "\cong"] & C^*(H) \arrow[r] \arrow[d, "\cong"] & 0 \\
0 \arrow[r] & C_0(H^* \times \mathbb{R}^\times) \arrow[r] & C_0(H^* \times \mathbb{R}^*) \arrow[r] & C_0(H^*) \arrow[r] & 0.
\end{tikzcd}$$

Observe that the bottom sequence is just

$$0 \to C_0(T^*X \smallsetminus H^*) \to C_0(T^*X) \to C_0(H^*) \to 0,$$

induced by the inclusion $H^* \subset T^*X$. Following our analysis of $C^*(G)$ we see that the KK-equivalence $I_H \sim C_0(H^* \times \mathbb{R}^\times)$ is a composition of the $*$-isomorphism

$$I_H \cong C^*(H \times \mathbb{R}^\times, s\omega) \cong C_0(\mathbb{R}) \otimes \mathcal{K}(V_-^{BF}) \oplus C_0(\mathbb{R}) \otimes \mathcal{K}(V_+^{BF})$$

with the Morita equivalence

$$\cdots \sim C_0(X \times \mathbb{R}) \oplus C_0(X \times \mathbb{R})$$

and, finally, the two Thom isomorphisms $\tau^- \oplus \tau^+$,

$$\cdots \sim C_0(H^{0,1} \times \mathbb{R}) \oplus C_0(H^{1,0} \times \mathbb{R}) \cong C_0(H^* \times \mathbb{R}^\times).$$

Here τ^+ and τ^- denote the Thom classes for the complex bundles $H^{1,0}$ and $H^{0,1}$ respectively.

Let us isolate the maps in K-theory that are relevant for our purposes. We summarize the conclusion of our analysis in the form of a proposition.

Proposition 2.6.1. *The following diagram commutes*

$$\begin{tikzcd}
K^0(T^*X \smallsetminus H^*) \arrow[r] & K^0(T^*X) \arrow[dd, "\Psi"', "\textit{Connes–Thom}"] \\
K^0(X \times \mathbb{R}^\times) \arrow[u, "\cup\tau^\pm", "\textit{Thom isom.}"'] & \\
K_0(I_H) \arrow[u, "\otimes V_\pm^{BF}", "\textit{Morita equiv.}"'] \arrow[r] & K^0(C^*(T_HX))
\end{tikzcd}$$

All vertical maps in the diagram are isomorphisms, and the horizontal maps (induced by inclusions) are surjective. The vertical arrows on the left are explicitly given by

$$\begin{tikzcd}
K^0(H^{0,1} \times \mathbb{R}) \oplus K^0(H^{1,0} \times \mathbb{R}) \arrow[r, "\cong"] & K^0(T^*X \smallsetminus H^*) \\
K^0(X \times \mathbb{R}) \oplus K^0(X \times \mathbb{R}) \arrow[u, "\cup\tau^- \oplus \cup\tau^+", "\textit{Thom isom.}"'] \arrow[r, "\cong"] & K^0(X \times \mathbb{R}^\times) \arrow[u] \\
K_0(C_0(\mathbb{R}) \otimes \mathcal{K}(V_-^{BF})) \ \oplus \ K_0(C_0(\mathbb{R}) \otimes \mathcal{K}(V_+^{BF})) \arrow[u, "\otimes V_-^{BF} \oplus \otimes V_+^{BF}", "\textit{Morita equiv.}"'] \arrow[r, "\cong"] & K_0(I_H) \arrow[u]
\end{tikzcd}$$

Proof. We only need to check the claim about surjectivity of the maps represented by the horizontal arrows. The inclusion of each connected component of $T^*X \smallsetminus H^*$ into T^*X is a homotopy equivalence. Therefore $K^0(T^*X \smallsetminus H^*) \to K^0(T^*X)$ is surjective. Since all vertical maps are isomorphisms, the same is true for the map $K_0(I_H) \to K^*(C^*(T_HX))$. □

2.7. **Poincaré duality.**

Definition 2.7.1. The Poincaré duality map

$$b : K_0(C^*(T_H X)) \to K_0^{top}(X)$$

is defined by choosing an arbitrary lift of an element in $K_0(C^*(T_H X))$ to an element in $K_0(I_H)$, followed by the composition of maps

$$K_0(I_H) \cong K^1(X) \oplus K^1(X) \overset{\cap[X^-]\oplus\cap[X^+]}{\longrightarrow} K_0^{top}(X) \oplus K_0^{top}(X) \to K_0^{top}(X).$$

From left to right, these maps are (1) Morita equivalence induced by the Bargmann-Fok Hilbert modules V_+^{BF} and V_-^{BF}, (2) Poincaré duality for the two natural Spin^c structures on X and (3) addition of K-homology classes.

Remark 2.7.2. Let us repeat here that the implicit isomorphism

$$K^0(X \times \mathbb{R}^\times) \cong K^0(X \times \mathbb{R}) \oplus K^0(X \times \mathbb{R}) \cong K^1(X) \oplus K^1(X)$$

is chosen such that the identifications $(-\infty, 0) \approx \mathbb{R}$ and $(0, \infty) \approx \mathbb{R}$ are given by the map $s \mapsto \log|s|$, i.e., each component of $\mathbb{R}^\times$ is oriented from 0 to $\pm\infty$.

Remark 2.7.3. The K-theory class of the symbol $[\sigma_H(P)] \in K_0(C^*(T_H X))$ only depends on the principal symbol of P in the Heisenberg calculus, and is uniquely defined by P, independent of arbitrary choices. In section 4 we will see that choosing a representative in $K_0(I_H)$ amounts to perturbing the distribution $\sigma_H(P) \in \mathcal{E}'(T_H X)$ by a function in $C_c^\infty(T_H X)$ in order to give it an extra property that is not determined by the principal symbol alone. The choice of such a perturbation is always possible, but not uniquely determined by P. However, the equivalence class of $b(\sigma_H(P))$ in K-homology does not depend on the choice of this perturbation, and so the map b is well-defined. The proof of this fact is implicit in the proof of the following theorem. We will also see that the second order operators $\Delta_H + i\gamma T$ *do* determine a canonical (or 'preferred') element in $K_0(I_H)$.

Lemma 2.7.4. *The following diagram commutes,*

$$\begin{tikzcd}
K^0(X \times \mathbb{R}) \arrow[r, "\cup\tau_+"] \arrow[d, "\cong"] & K^0(T^*X) \arrow[d, "c"] \\
K^1(X) \arrow[r, "\cap[X^+]"'] & K_0^{top}(X)
\end{tikzcd}$$

Here the left vertical arrow is the suspension isomorphism in K-theory; the upper horizontal arrow is the Thom isomorphism resulting from the direct sum decomposition $T^*X = H^{1,0} \oplus \underline{\mathbb{R}}$; the lower horizontal arrow is Poincaré duality on the Spin^c manifold X, i.e., is cap product with the K-homology fundamental cycle $[X^+]$; and the right vertical arrow is the "clutching construction", i.e., cap product with the K-homology fundamental cycle of the Spin^c manifold T^*X.

Proof. Commutativity of the diagram follows from commutativity of the two triangles that appear if we introduce the diagonal arrow which is the Thom isomorphism for the Spin^c vector bundle T^*X on X,

$$\begin{tikzcd}
K^0(X\times\mathbb{R}) \arrow[r, "\cup\tau_+"] & K^0(T^*X) \arrow[d, "c"] \\
K^1(X) \arrow[u, "\cong"] \arrow[ur] \arrow[r, "\cap[X^+]"'] & K_0^{top}(X)
\end{tikzcd}$$

Commutativity of the two triangles is standard algebraic topology, which we now briefly indicate. The upper triangle commutes because the Thom class for $T^*X = H^{1,0}\oplus\underline{\mathbb{R}}$ is the product of the Thom classes used for the left vertical arrow and the upper horizontal arrow. For commutativity of the lower triangle, use the following notation: $\beta\in KK^1(C(X),C_0(T^*X))$ is the Thom class for T^*X, $\alpha\in KK^1(C_0(T^*X),C(X))$ is the KK-element given by the family of Dirac operators for the fibers of T^*X. The Dirac-dual Dirac identity in this context is the assertion that the Kasparov product $\beta\#\alpha$ is the unit element of the ring $KK^0(C(X),C(X))$,

$$\beta\#\alpha = 1\in KK^0(C(X),C(X)).$$

The total space of T^*X is itself a (non-compact) Spin^c manifold, and its K-homology fundamental class $[T^*X]$ is the Kasparov product of α and $[X^+]$,

$$[T^*X] = \alpha\#[X^+]\in KK^0(C_0(T^*X),\mathbb{C}).$$

It now follows that

$$\beta\#[T^*X] = \beta\#\alpha\#[X^+] = [X^+].$$

which gives commutativity of the lower triangle. □

Theorem 2.7.5. *The following diagram commutes*

$$\begin{tikzcd}
K^0(T^*X) \arrow[r, "\Psi"] \arrow[d, "c"] & K_0(C^*(T_HX)) \arrow[dl, "b"] \\
K_0^{top}(X) &
\end{tikzcd}$$

where b is as in Definition 2.7.1, and c is the "clutching construction" introduced in [6, section 22].

Proof. If we identify $X\times(0,\infty)\cong X\times\mathbb{R}$, then Lemma 2.7.4 gives commutativity of

$$\begin{tikzcd}
K^0(X\times(0,\infty)) \arrow[r, "\cup\tau_+"] \arrow[d, "\cong"] & K^0(T^*X) \arrow[d, "c"] \\
K^1(X) \arrow[r, "\cap[X^+]"'] & K_0^{top}(X)
\end{tikzcd}$$

Similarly, for the conjugate Spin^c structure on X we have

$$\begin{tikzcd}
K^0(X\times(-\infty,0)) \arrow[r, "\cup\tau_-"] \arrow[d, "\cong"] & K^0(T^*X) \arrow[d, "c"] \\
K^1(X) \arrow[r, "\cap[X^-]"'] & K_0^{top}(X)
\end{tikzcd}$$

The *sign* of the isomorphism $K^0(X\times(0,\infty))\cong K^1(X)$ depends on the orientation of $(0,\infty)$, i.e., on the orientation of the normal bundle $N = X\times\mathbb{R}$, which in turn depended on the choice of contact form. In order to make the second diagram commute we must orient $(-\infty,0)$ from 0 to $-\infty$. This is because the reversed orientation of the normal bundle N is built into the definition of the fundamental cycle $[X^-]$, and also affects the choice of the isomorphism

$$K^0(X\times(-\infty,0))\cong K^0(X\times\mathbb{R})\cong K^1(X).$$

Combining the two components of $X\times\mathbb{R}^\times$, we obtain a single diagram

$$\begin{tikzcd}
K^0(X\times(-\infty,0))\oplus K^0(X\times(0,+\infty)) \arrow[r, "\cup\tau^-\oplus\cup\tau^+"] \arrow[d, "\cong"] & K^0(T^*X\smallsetminus H^*) \arrow[r] & K^0(T^*X) \arrow[d, "c"] \\
K^1(X)\oplus K^1(X) \arrow[r, "\cap[X^-]\oplus\cap[X^+]"'] & K_0^{top}(X)\oplus K_0^{top}(X) \arrow[r] & K_0^{top}(X)
\end{tikzcd}$$

Comparing this diagram with Corollary 2.6.1 proves the proposition.

□

3. The index theorem as a commutative triangle

The symbol $\sigma_H(P)$ of a hypoelliptic operator in the Heisenberg calculus naturally determines an element in $K_0(C^*(T_HX))$. In this section we prove that the Poincaré dual $b(\sigma_H(P))\in K_0^{top}(X)$ of the Heisenberg symbol is the desired K-cycle $\mu^{-1}(P)$. The main result of this section is Theorem 3.5.1 which asserts that there is commutativity in the triangle

$$\begin{tikzcd}
& K_0(C^*(T_HX)) \arrow[dl, "b"'] \arrow[dr, "\mathrm{Op_H}"] & \\
K_0(X) \arrow[rr, "\mu"] & & KK_0(C(X),\mathbb{C})
\end{tikzcd}$$

For convenience of the reader, we start with a brief sketch of the main features of the Heisenberg calculus. References are [10, 17, 23].

3.1. **Differential operators in the Heisenberg calculus.** Consider a differential operator P on a smooth manifold X, given in local coordinates by an expression

$$P=\sum_{|\alpha|\le d}a_\alpha\partial^\alpha,\qquad \alpha=(\alpha_1,\ldots,\alpha_n),\ |\alpha|=\sum\alpha_j,\ \partial^\alpha=\Pi(\partial/\partial x_j)^{\alpha_j},$$

where the coefficients a_α are smooth functions. The highest order part of P is not well-defined as a differential operator on X. The algebra of differential operators is only filtered, not graded. But the highest order part at a point $x\in X$,

$$P_x=\sum_{|\alpha|=d}a_\alpha(x)\partial^\alpha,$$

can be interpreted as a constant coefficient operator on the tangent fiber T_xX. As such it is well-defined and independent of coordinate choices.

The root of the *Heisenberg calculus* is a simple but important idea, proposed by Gerald Folland and Elias Stein in [18], that it is useful for the analysis of certain hypoelliptic operators to equip the algebra of differential operators with an alternative filtration. The filtration on the algebra of differential operators proposed by Folland and Stein is generated by a filtration on the Lie algebra of vector fields, defined as follows: All vector fields in the direction of the contact hyperplane bundle H have order one, as usual, but *any* vector field that is not everywhere tangent to H is given order *two*. For example, a sublaplacian Δ_H has order two, as in the classical calculus. But in the Heisenberg calculus the Reeb vector field T is also a second order operator.

What sort of object is the highest order part of a differential operator if we adopt this Heisenberg filtration? Abstractly, for any filtered algebra, the notion of 'highest order part' refers to an element in the associated graded algebra. Observe that in the associated graded algebra smooth functions f commute with all vector fields W, because the commutator $[W, f] = W.f$ is of order zero. This implies that elements in the graded algebra can be *localized* at points $x \in X$. Therefore, just as in the classical case, the highest order part of a differential operator P will consist of a family of operators P_x parametrized by $x \in X$. But they are not exacty constant coefficient operators here. Instead, they are translation invariant operators for a certain nilpotent group structure on T_xX.

In all that follows, X denotes a closed contact manifold with a contact hyperplane bundle $H \subset TX$. We denote by $N = TX/H$ the quotient line bundle, and assume that there exists a global contact form θ, so that we can identify $N = X \times \mathbb{R}$. Also, the Reeb vector field provides us with a section $N \subset TX$, so that we can identify $T_xX = H_x \times \mathbb{R}$, and that $d\theta$ restricted to H_x is a symplectic form. The tangent space $G_x = H_x \times \mathbb{R}$ is then a Heisenberg group with group operation

$$(v, t) \cdot (v', t') = (v + v', t + t' - \frac{1}{2} d\theta(v, v')), \qquad v, v' \in H_x,\ t, t' \in \mathbb{R}.$$

Algebraically, the smooth groupoid T_HX is the disjoint union of Heisenberg groups

$$T_HX = \bigsqcup_{x \in X} G_x.$$

The highest order part $\{P_x, x \in X\}$ of a differential operator P in the Heisenberg calculus can be interpreted as a smooth family of translation invariant operators P_x on the Heisenberg groups $G_x = H_x \times \mathbb{R} \cong T_xX$, or, equivalently, a right invariant differential operator on the Lie groupoid T_HX.

For example, for the second order operators $P_\gamma = \Delta_H + i\gamma T$, freezing coefficients at a point $x \in X$ results in

$$P_x = -\sum_{j=1}^{2n} W_j(x)^2 + i\gamma(x)T(x).$$

This formal polynomial in the tangent vectors $W_j(x)$, $T(x) \in T_xX$ should be interpreted as an element in the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_x)$ of the Heisenberg Lie algebra $\mathfrak{g}_x = H_x \oplus \mathbb{R}$, or, equivalently, as an invariant differential operator on the group G_x. In other words, the vector $W_j(x)$ should *not* be identified with a vector field on T_xX that is translation invariant for the usual vector space structure on T_xX, but rather with a vector field on T_xX that is translation invariant for the Heisenberg group structure on T_xX.

3.2. **Heisenberg pseudodifferential calculus.** When we restrict attention to differential operators the Heisenberg calculus is fairly straightforward. Constructing the corresponding $\mathbb{Z}$-filtered algebra of Heisenberg *pseudodifferential* operators requires more work (see [10, 17, 23]). In this section we sketch one possible approach.

Just like the classical pseudodifferential calculus, the Heisenberg algebra consists of pseudolocal continuous linear operators

$$P : C^\infty(X) \to C^\infty(X).$$

In other words, they are operators with a Schwartz kernel $k(x,y)$ that is a smooth function off the diagonal in $X\times X$. The Heisenberg calculus (as well as its filtration) is defined by asymptotic expansions of the singularity of k in the direction transversal to the diagonal.

Choose a connection ∇ on TX that preserves the distribution H (this is an important technical condition). Consider the exponential map $\exp^{\nabla}$ associated to this connection ∇,

$$h : T_H X \to X\times X : (x,v) \mapsto (\exp_x^{\nabla}(v), x).$$

The map h is a local diffeomorphism of a neighborhood of the zero section of $T_H X$ with a neighborhood of the diagonal in $X\times X$. We pull back the distribution k to the groupoid $T_H X$ by the map h, and then chop it by a cut-off function. Let ϕ be an arbitrary smooth function on $T_H X$ that is compactly supported and equals 1 in a neighborhood of the zero section. At a point $x\in X$ let k_x be the distribution on the Heisenberg group $G_x = T_x X$ defined by

$$k_x(v) = \phi(x,v)\cdot k(h(x,-v)), \qquad v\in T_x X.$$

The smooth family of distributions $\{k_x, x\in X\}$ defines a compactly supported distribution on $T_H X$. The action of the operator P with kernel k is approximated (in a neighborhood of the point $x\in X$) by convolution with the compactly supported distribution k_x on the Heisenberg group G_x, in a sense that can be made precise. This was the basic idea introduced by Folland and Stein in [18].

Because the kernel k is smooth off the diagonal, each distribution k_x is regular, i.e., it is a smooth function when restricted to $G_x \setminus \{0\}$. We say that P is a pseudodifferential operator of order d if each distribution k_x has an asymptotic expansion near $0\in G_x$,

$$k_x \sim k_x^0 + k_x^1 + k_x^2 + \cdots$$

The asymptotic expansion should be interpreted in the usual way: the remainder $k - \sum_{j=0}^N k^j$ becomes more regular as N increases, and the entire expansion determines k_x modulo compactly supported smooth functions $C_c^\infty(T_H X)$. The defining feature of the Heisenberg calculus is that the terms k_x^j in this expansion must be homogeneous with respect to the 'parabolic' dilation structure of the Heisenberg group $G_x = T_x X = H_x \times \mathbb{R}$,

$$\delta_s : H_x\times\mathbb{R} \to H_x\times\mathbb{R} : (v,t)\mapsto (sv, s^2 t), \qquad s>0.$$

Formally, the term k_x^j in the expansion must satisfy

$$k_x^j(\delta_s v) = s^{j-(2n+2)}\, k_x^j(v).$$

This notion of homogeneity based on the dilations δ_s corresponds precisely to the Heisenberg filtration that assigns order one to vector fields tangent to H and order two to vector fields transveral to H. But we now obtain a filtered algebra of pseudodifferential operators that differs from the classical pseudodifferential algebra.

The existence of an asymptotic expansion is independent of the choice of exponential map $\exp^{\nabla}$ and cut-off function ϕ, as long as the exponential map satisfies the technical condition that it preserves sections in H. We denote the operator of convolution with the distribution k_x by P_x, and we write

$$\sigma_H(P) = \{P_x, x\in X\}.$$

We regard $\sigma_H(P)$ as an element in the convolution algebra $\mathcal{E}'(T_H X)$ of compactly supported distributions on the groupoid $T_H X$, and we can think of it as the 'full symbol' of P. While the value of $\sigma_H(P)$ depends on the choice of exponential $\exp^{\nabla}$ and cut-off ϕ, the highest order part $\sigma_H^d(P) = \{k_x^0, x\in X\}$ in the asymptotic expansion of $\sigma_H(P)$ is invariantly defined as a smooth family of convolution operators on $T_H X$, independent of $\exp^{\nabla}, \phi$.

The collection of all Heisenberg pseudodifferential operators on X is a $\mathbb{Z}$-filtered algebra: if P, Q are Heisenberg pseudodifferential operators of order a, b respectively, then PQ is a Heisenberg pseudodifferential operator of order $a+b$. Moreover, the leading term in the asymptotic expansion of $\sigma_H(PQ)$ is obtained by convolution of the leading terms in the expansions of $\sigma_H(P)$ and $\sigma_H(Q)$, i.e.,

$$\sigma_H^{a+b}(PQ) = \sigma_H^a(P) * \sigma_H^b(Q).$$

As usual, if the leading term $\sigma_H^d(P)$ of an order d symbol $\sigma_H(P)$ is invertible (in the convolution algebra), then we can derive an asymptotic expansion for an inverse of the full symbol $\sigma_H(P)$ modulo $C_c^\infty(T_HX)$. This implies that P itself is invertible in the Heisenberg algebra modulo smoothing operators, and hence a hypoelliptic Fredholm operator on X. It is precisely for such operators that we prove our index formula.

3.3. **The Heisenberg symbol in K-theory.** In [24, Definition 15] we constructed a K-theoretic symbol in $K_0(C^*(T_HX))$ for a hypoelliptic differential operator P in the Heisenberg calculus. As we have seen, the principal symbol $\sigma_H^d(P)$ is then a family of differential operators P_x on G_x, where each P_x is obtained from P by a simple procedure of 'freezing coefficients' at $x \in X$. The result is independent of a choice of exponential map $T_HX \to X \times X$ or cut-off function ϕ. In fact, the construction given in [24] makes no reference to the Heisenberg pseudodifferential calculus at all. In [25] we gave an alternative definition that works for order zero operators in the Heisenberg calculus.

Perhaps the easiest way to define the K-theory element

$$[\sigma_H(P)] \in K_0(C^*(T_HX))$$

is to follow the general ideas set out by Connes in [16, II.9.α]. The resulting construction works for arbitrary hypoelliptic operators in the Heisenberg algebra.

Compactly supported smooth functions $C_c^\infty(T_HX)$ form a two-sided ideal in the convolution algebra $\mathcal{E}'(T_HX)$ of compactly supported distributions on the groupoid T_HX. The asymptotic expansion for products of full symbols in the Heisenberg calculus implies that if the principal symbol of an operator P is invertible, then the *full* symbol $\sigma_H(P)$ has an inverse modulo the ideal $C_c^\infty(T_HX)$. By a very general argument explained by Connes, this implies that $\sigma_H(P)$ has an 'index' in the K-theory group $K_0(C^*(T_HX))$. First, the full symbol $\sigma_H(P)$ determines an element in *algebraic* relative K-theory,

$$[\sigma_H(P)] \in K_0(\mathcal{E}'(T_HX), C_c^\infty(T_HX)).$$

By excision we have

$$K_0(\mathcal{E}'(T_HX), C_c^\infty(T_HX)) \cong K_0(C_c^\infty(T_HX)),$$

and, finally, the inclusion of rings $C_c^\infty(T_HX) \subset C^*(T_HX)$ maps the element $[\sigma_H(P)]$ to the K-theory group $K_0(C^*(T_HX))$ (which is the same in algebraic and C^*-algebraic K-theory). To make this more explicit, if Q is a parametrix for P (an inverse modulo smoothing operators), then the formal difference of idempotents

$$[\sigma_H(P)] = [e] \ominus [e_0] \in K_0(C^*(T_HX))$$

is given by

$$e = \begin{pmatrix} S_0^2 & S_0\sigma_H(Q) \\ S_1(1+S_1)\sigma_H(P) & 1-S_1^2 \end{pmatrix}, \; e_0 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix},$$

where $S_0 = 1 - \sigma_H(Q)\sigma_H(P)$, $S_1 = 1 - \sigma_H(P)\sigma_H(Q)$. (See [16, II.9.$\alpha$] for a detailed discussion of the general construction of a K-theoretic index of an element in a ring that is invertible modulo an ideal.)

3.4. **The "Choose an operator" maps.** In this section we prove the analog of the Poincaré duality Theorem 2.7.5 for analytic K-homology $KK(C(X),\mathbb{C})$. The proof of this theorem closely follows the steps of [24, section 3.7], generalizing everything from a statement in K-theory to a statement about the functor $KK(C(X),-)$. We refer the reader to [24] for details of the proof in K-theory, and indicate here how the argument can be stengthened to prove Theorem 3.4.1.

The Poincaré duality map

$$\mathrm{Op}_e : K^0(T^*X) \to KK^0(C(X),\mathbb{C})$$

is defined as follows. Given $\sigma \in K^0(T^*X)$, choose an elliptic zero order pseudodifferential operator P on X such that in $K^0(T^*X)$,

$$[\sigma(P)] = \sigma.$$

Then $\mathrm{Op}_e(\sigma)$ is the element in $KK^0(C(X),\mathbb{C})$ determined by P. The non-trivial point here is that if two elliptic pseudodifferential operators P,Q have $[\sigma(P)] = [\sigma(Q)]$ in $K^0(T^*X)$, then P,Q determine the same element in $KK^0(C(X),\mathbb{C})$—i.e., a homotopy of symbols can be lifted to a homotopy of operators. (See [2], [6, section 23].) Instead of attempting a direct construction for the Heisenberg pseudodifferential calculus, we will, in the proof below, construct a map

$$\mathrm{Op}_H : K_0(C^*(T_HX)) \to KK(C(X),\mathbb{C}),$$

and then prove a posteriori that if P is an order zero operator in the Heisenberg calculus with principal symbol $\sigma_H(P)$, then

$$\mathrm{Op}_H([\sigma_H(P)]) = [P].$$

This justifies referring to the map Op_H as the "choose a hypoelliptic operator" map for the Heisenberg calculus.

Theorem 3.4.1. *There exists a map*

$$\mathrm{Op}_H : K_0(C^*(T_HX)) \to KK^0(C(X),\mathbb{C}),$$

such that (i) if P is an order zero operator in the Heisenberg algebra with invertible principal symbol $\sigma_H(P)$ then $\mathrm{Op}_H(\sigma_H(P)) = [P]$ in $KK^0(C(X),\mathbb{C})$, and (ii) the following diagram commutes,

$$\begin{tikzcd}
K^0(T^*X) \arrow[r, "\Psi"] \arrow[d, "\mathrm{Op_e}"] & K_0(C^*(T_HX)) \arrow[dl, "\mathrm{Op_H}"] \\
KK^0(C(X),\mathbb{C}) &
\end{tikzcd}$$

Proof. The proof is a strengthening of the groupoid arguments developed in [24]. The main difference is that we need to replace K-theory $KK(\mathbb{C},-)$ by the functor $KK(C(X),-)$. What makes this possible is that all C^*-algebras that play a role in our argument are of Type I, and hence nuclear. Therefore all ideals are semi-split (by [13]), and since $C(X)$ is separable (X is a compact manifold) the $KK(C(X),-)$ functor has 6-term exact sequences [11, Thm. 19.5.7]. As a result, the quotients by various contractible ideals that play a role in the tangent groupoid argument induce isomorphisms not only in $K_0(-) \cong KK(\mathbb{C},-)$, but also in $KK(C(X),-)$.

We briefly review the argument and indicate how it can be strengthened. For a differential Heisenberg operator P with invertible principal symbol we constructed in [24, section 3.6] an element in the K-theory of the parabolic tangent groupoid,

$$[\mathbb{P}] \in K_0(C^*(\mathbb{T}_H X)).$$

The 'parabolic tangent groupoid' $\mathbb{T}_H X$ is algebraically the disjoint union of $T_H X$ and the family of pair groupoids $X \times X \times (0,1]$, similar to Connes' tangent groupoid. The 'blow-up' of the diagonal in $X \times X$ is defined using the parabolic dilations δ_s in the Heisenberg groups G_x [24, section 3]. The class $[\mathbb{P}]$ combines the principal Heisenberg symbol of P with P itself into a single K-theory element for the tangent groupoid.

We must modify this construction and apply it to order zero operators. This is actually a considerable simplification, especially if we construct $[\mathbb{P}]$ in KK-theory. Let $\mathcal{E} = C^*(\mathbb{T}_H X) \oplus C^*(\mathbb{T}_H X)$ be the obvious $\mathbb{Z}_2$-graded Hilbert module over $C^*(\mathbb{T}_H X)$. The adjointable operator

$$\mathbb{P} \in \mathcal{L}(\mathcal{E}) \cong M(C^*(\mathbb{T}_H X)) \otimes M_2(\mathbb{C})$$

restricts to the principal symbol

$$\begin{pmatrix} 0 & P_x^* \\ P_x & 0 \end{pmatrix} \in M(C^*(G_x)) \otimes M_2(\mathbb{C})$$

for the fiber at $s = 0$ and $x \in X$, and to

$$\begin{pmatrix} 0 & P^* \\ P & 0 \end{pmatrix} \in \mathcal{L}(L^2(M) \oplus L^2(M)) \cong M(C^*(X \times X)) \otimes M_2(\mathbb{C})$$

for all other values $s > 0$. The principal symbol is, by definition, the highest order part in an asymptotic expansion of the operator kernel of P, and these asymptotics agree precisely with the parabolic 'blow-up' of the diagonal in $X \times X$ that defines $\mathbb{T}_H X$.

What we did not point out in [24] is that $(\mathcal{E}, \mathbb{P})$ actually defines an element in

$$[\mathbb{P}] \in KK(C(X), C^*(\mathbb{T}_H X)).$$

There is an obvious diagonal representation

$$\phi : C(X) \to \mathcal{L}(\mathcal{E}) = M(C^*(\mathbb{T}_H X)) \otimes M_2(\mathbb{C}).$$

At $s = 0$ it restricts to the identification of $C(X)$ with the center of $M(C^*(T_H X))$, while at $s > 0$ we have the representation of continuous functions on X as multiplication operators in $M(C^*(X \times X)) \cong \mathcal{L}(L^2(X))$. One easily verifies compactness of commutators $[\mathbb{P}, \phi(f)] \in \mathcal{K}(\mathcal{E})$ for continuous functions $f \in C(X)$.

Just as in [24], the restriction map at $s = 0$ has a contractible ideal and therefore induces an isomorphism

$$e_0 : KK(C(X), C^*(\mathbb{T}_H X)) \cong KK(C(X), C^*(T_H X)).$$

Observe that, by construction of the Fredholm module $[\mathbb{P}]$, we have $e_0([\mathbb{P}]) = [\sigma_H(P)]$.

Restriction to the fiber at $s = 1$ induces a map

$$e_1 : KK(C(X), C^*(\mathbb{T}_H X)) \to KK(C(X), C^*(X \times X)) \cong KK(C(X), \mathbb{C}).$$

Again, by construction, $e_1([\mathbb{P}]) = [P]$. It follows that the combined map

$$e_1 \circ e_0^{-1} : KK(C(X), C^*(T_H X)) \to KK(C(X), \mathbb{C}),$$

is the "choose a hypoelliptic operator" map,

$$e_1 \circ e_0^{-1}([\sigma_H(P)]) = [P].$$

The exact same argument, using the tangent groupoid $\mathbb{T}X = TX \sqcup X \times X \times (0,1]$ of Connes, works for elliptic order zero operators.

Having proven that the "choose an operator" map is induced by the appropriate tangent groupoid, we can now show that these two maps commute with the isomorphism Ψ. The argument from [24, section 3.8] involving the *adiabatic groupoid* of the parabolic tangent groupoid $\mathbb{T}_H X$ (a deformation of a deformation) applies, without essential change, to show that the diagram

$$\begin{tikzcd} KK(C(X), C_0(T^*X)) \arrow[r, "\Psi"] \arrow[d] & KK(C(X), C^*(T_H X)) \arrow[dl] \\ KK((C(X), \mathbb{C}) & \end{tikzcd}$$

commutes.

If $\mathcal{H}_A$ denotes the standard countably generated free Hilbert A-module, then any $*$-homomorphism $\phi : C(X) \to Z(M(A))$ induces, in the obvious way, a map $\tilde{\phi} : C(X) \to Z(\mathcal{L}(\mathcal{H}_A))$. Thus, for a $C(X)$-C^*-algebra A there is a natural homomorphism

$$\alpha_X : KK(\mathbb{C}, A) \to KK(C(X), A).$$

The map α_X is a natural transformation from the $KK(\mathbb{C}, -)$ functor to the $KK(C(X), -)$ functor on the category of $C(X)$-C^*-algebras. In fact, it is a natural transformation for the KK^X-category.

Therefore, since the isomorphism Ψ is implemented by a KK^X-element

$$\Psi \in KK^X(C_0(T^*X), C^*(T_H X))$$

we have a commutative diagram

$$\begin{tikzcd} KK(\mathbb{C}, C_0(T^*X)) \arrow[r, "\Psi"] \arrow[d, "\alpha_X"'] & KK(\mathbb{C}, C^*(T_H X)) \arrow[d, "\alpha_X"] \\ KK(C(X), C_0(T^*X)) \arrow[r, "\Psi"] & KK(C(X), C^*(T_H X)) \end{tikzcd}$$

Composing this with our result above yields the proposition.

□

3.5. The hypoelliptic index theorem in K-homology.

The Atiyah-Singer theorem for elliptic operators amounts to commutativity of the triangle [6, Thm. 23.1],

$$\begin{tikzcd} & K^0(T^*X) \arrow[dl, "c"'] \arrow[dr, "\mathrm{Op_e}"] & \\ K_0(X) \arrow[rr, "\mu"] & & KK_0(C(X), \mathbb{C}). \end{tikzcd}$$

We obtain an analogous theorem for the Heisenberg calculus.

Theorem 3.5.1. *Let (X, H) be a closed contact manifold. Then there is commutativity in the diagram*

$$\begin{tikzcd} & K_0(C^*(T_H X)) \arrow[dl, "b"'] \arrow[dr, "\mathrm{Op_H}"] & \\ K_0(X) \arrow[rr, "\mu"] & & KK_0(C(X), \mathbb{C}) \end{tikzcd}$$

Proof. The theorem is implied by commutativity of the three smaller triangles in the diagram,

$$\begin{tikzcd}
& K_0(C^*(T_HX)) \arrow[ddl, bend right, "b"'] \arrow[ddr, bend left, "\mathrm{Op_H}"] & \\
& K^0(T^*X) \arrow[u, "\Psi"] \arrow[dl, "c"] \arrow[dr, "\mathrm{Op_e}"'] & \\
K_0^{top}(X) \arrow[rr, "\mu"'] & & KK^0(C(X),\mathbb{C})
\end{tikzcd}$$

□

Modulo an explicit computation of the Poincaré dual $b(\sigma_H(P))$ of the Heisenberg symbol of a hypoelliptic operator, Theorem 3.5.1 solves the general index problem for hypoelliptic operators in the Heisenberg calculus. We will now apply Theorem 3.5.1 to find explicit (M,E,φ) cycles for certain examples of hypoelliptic second order differential operators (in the Heisenberg calculus) on X.

4. Computation of the K-cycle $b(\sigma_H(P))$

We now apply Theorem 3.5.1 to compute the geometric K-cycle $b(\sigma_H(P_\gamma))$ for the second order differential operators

$$P_\gamma = \Delta_H + i\gamma T.$$

Also, we answer a question posed by R. Ponge, and indicate how Louis Boutet de Monvel's formula for the index of Toeplitz operators is a corollary of Theorem 3.5.1. Finally we consider the case of odd-dimensional spheres.

4.1. **Perturbing the symbol.**

We wish to compute the image of the K-theory element associated to the Heisenberg symbol $\sigma_H(P_\gamma)$ of P_γ under the Poincaré duality map of Definition 2.7.1,

$$b : K_0(C^*(T_HX)) \to K_0(X)$$

The first step is to lift $[\sigma_H(P)]$ to a class in $K_0(I_H)$. The precise details of this lift turn out to be irrelevant. Recall that the operator P determines the distribution $\sigma_H(P)$ only up to a perturbation by $C_c^\infty(T_HX)$. And one can always choose a perturbation of $\sigma_H(P)$ by a $C_c^\infty(T_HX)$ function that defines an element in $K_0(I_H)$.

The full symbol $\sigma_H(P) = \{P_x, x \in X\}$ is an element in the convolution algebra $\mathcal{E}'(T_HX)$ of compactly supported distributions on T_HX. If P is hypoelliptic, then $\sigma_H(P)$ is invertible modulo $C_c^\infty(T_HX)$. Consider the algebra homomorphism

$$\pi_0 : \mathcal{E}'(T_HX) \to C^\infty(H^*),$$

defined as the composition of Fourier transform in the transversal $\mathbb{R}$-variable of $T_HX = H \times \mathbb{R}$ and restriction to $s = 0$ in $\mathbb{R}^*$. If P is an order zero operator then the function $\pi_0(\sigma_H(P))$ extends to a smooth function on the radial compactification

$$DH^* = H^* \sqcup SH^*.$$

The restriction of $\pi_0(\sigma_H(P))$ to the sphere bundle SH^* is the *equatorial symbol* of Epstein and Melrose [17]. The equatorial symbol is invertible if P is hypoelliptic. In fact, if P is hypoelliptic then the function $\pi_0(\sigma_H(P))$ is invertible modulo $C_c^\infty(H^*)$.

Now consider the exact sequence in K-theory

$$K^0(T^*X \setminus H^*) \to K^0(T^*X) \to K^0(H^*).$$

The first map is surjective, and so the second map is trivial. Therefore the same is true for the isomorphic sequence

$$K_0(I_H) \to K_0(C^*(T_H X)) \xrightarrow{\pi_0} K_0(C_0(H^*)).$$

We see that $\pi_0(\sigma_H(P))$ defines a trivial element in $K_0(C^*(H))$. It follows that $\pi_0(\sigma_H(P))$ can be perturbed by a function in $C_c^\infty(H^*)$ (possibly a matrix valued function) to obtain an invertible function on all of DH^*. And so the distribution $\sigma_H(P)$ can be perturbed by a function in $C_c^\infty(T_H X)$ to obtain a new element in $\mathcal{E}'(T_H X)$ for which $\pi_0(\sigma_H(P))$ is invertible. The resulting perturbation defines an element in $K_0(I_H)$.

4.2. Suspension in K-theory.

The first step in Definition 2.7.1 is the isomorphism

$$K_0(I_H) \cong K^0(X \times \mathbb{R}^\times) \cong K^1(X) \oplus K^1(X),$$

which is a composition of Morita equivalence and suspension. We may reverse the order of the Morita equivalence and the suspension isomorphisms, and proceed via

$$K_0(I_H) \cong K_1(\mathcal{K}(V_-^{BF})) \oplus K_1(\mathcal{K}(V_+^{BF})) \cong K^1(X) \oplus K^1(X).$$

We will need an explicit formula for the suspension isomorphism. Since we could not find a reference for the following simple K-theory lemma, we provide a proof here.

Lemma 4.2.1. *Let B be a separable C^*-algebra. Let $u \in M_n(B)^+$ be a unitary matrix that represents a class $[u] \in K_1(B)$, and let θ denote the suspension isomorphism,*

$$\theta : K_1(B) \to K_0(C_0(0,1) \otimes B).$$

If $F(t)$, $t \in [0,1]$ is the norm continuous family that linearly interpolates between $F(0) = 1$ and $F(1) = u$,

$$F(t) = 1 - t + tu \in M_n(B)^+,$$

then $F \in C_0([0,1], M_n(B))^+$ is invertible modulo $C_0((0,1), M_n(B))$, and defines a relative class

$$[F] \in K_0(C_0(0,1) \otimes B).$$

Then $[F] = \theta([u])$.

Proof. We will prove this by making use of the flexibility of KK-theory. For ease of notation, let us write $A = C_0((0,1), B)$. Since $M_n(B)^+ \subset M_n(M(A)) = \mathcal{L}(A^n)$, we can think of F as an adjointable operator on the Hilbert A-module A^n. The fact that $F_0 = 1$, $F_1 = u$ and $F_t = 1$ modulo $M_n(B)$ for all $t \in [0,1]$ implies that F is invertible modulo compact operators $\mathcal{K}(A^n) = C_0((0,1), M_n(B))$. In other words, the relative K-theory class defined by F can also be represented as a Kasparov module

$$[F : A^n \to A^n] \in KK(\mathbb{C}, A).$$

We will prove that this Fredholm module corresponds to $\theta([u])$ under the standard identification $KK(\mathbb{C}, A) \cong K_0(A)$.

Choose a continous family of unitaries $z(t) \in M_{2n}(B^+), t \in [0,1]$ such that

$$z_0 = \begin{pmatrix} 1_n & 0 \\ 0 & 1_n \end{pmatrix},\ z_1 = \begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix}.$$

Let

$$p_n = \begin{pmatrix} 1_n & 0 \\ 0 & 0 \end{pmatrix},$$

and let e denote the continuous family of projections

$$e_t = z_t p_n z_t^{-1}.$$

Observe that $e_0 = e_1 = p_n$. Then the proof of [Blackadar, Theorem 8.2.2] implies that

$$\theta([u]) = [e] - [p_n] \in K_0(C_0((0,1),B)).$$

Consider the family of partial isometries

$$G(t) = z(t)p_n.$$

Observe that $G(0) = p_n$, $G(1) = u$ and that $G(1) = p_n$ modulo $\mathcal{K}(A^{2n})$, while

$$1 - G^*G = 1 - p_n,\ 1 - GG^* = 1 - e.$$

Let $\phi : \mathbb{C} \to \mathcal{L}(B^{2n})$ denote the homomorphism with $\phi(1) = p_n$. With this choice of a nonunital map ϕ the operator G satisfies the Fredholm property, and we obtain a Kasparov module

$$[G : A^{2n} \to A^{2n}, \phi] \in KK(\mathbb{C}, A).$$

Moreover, the module $[G : A^{2n} \to A^{2n}, \phi]$ is homotopic to $[F : A^n \to A^n, p_n\phi p_n]$, where $p_n\phi p_n : \mathbb{C} \to \mathcal{L}(A^n)$ is just the standard unital map. But unlike F the operator G has closed range, and we can take its index

$$\operatorname{Index} G = [1 - G * G] - [1 - GG^*] = [e] - [p_n] = \theta([u]).$$

It follows that $[G]$ corresponds to $\theta([u])$, and therefore so does $[F]$. □

4.3. **Second order scalar operators.** We have computed the Fredholm index of the hypoelliptic operators $P_\gamma = \Delta_H + i\gamma T$ in [25] (see the example following [25, Prop. 7]). We can now identify the full K-cycle. Our K-cycle for P_γ contains new information that cannot be inferred from the formula in [25].

Proposition 4.3.1. *Let* $\tilde{P}_\gamma = P_\gamma(1 + P_\gamma^* P_\gamma)^{-1/2}$. *Then*

$$b(\sigma_H(P_\gamma)) = [\pi_-(\sigma_H(\tilde{P}_\gamma))] \cap [X^-] \ + \ [\pi_+(\sigma_H(\tilde{P}_\gamma))] \cap [X^+].$$

Here π_+ *denotes the Bargmann-Fok representation of the groupoid* $T_H X$ *on the Hilbert module* V_+^{BF}, *while* π_- *denotes the dual representation on* V_-^{BF}.

Proof. Observe that the equatorial symbol of the order zero operator $\tilde{P}_\gamma$ is identically equal to 1. Therefore there exists a $C_c^\infty(T_H X)$ perturbation of $\sigma_H(\tilde{P}_\gamma)$ such that the function $\pi_0(\sigma_H(\tilde{P}_\gamma))$ becomes identically 1 on all of DH^*.

Let $B = \mathcal{K}(V_-^{BF}) \oplus \mathcal{K}(V_+^{BF})$. Then

$$I_H \cong C_0((0,\infty), B).$$

Let u be the invertible element

$$u = \pi_-(\sigma_H(\tilde{P}_\gamma)) \oplus \pi_+(\sigma_H(\tilde{P}_\gamma)) \in B^+.$$

The difference between $\sigma_H(\tilde{P}_\gamma)$ and $1 - t + tu$ is an element in $C_0((0,1),B)$, so that $\sigma_H(\tilde{P}_\gamma)$ and $1 - t + tu$ represent the same element in $K_0(I_H)$. Lemma 4.2.1 now implies that the K-theory element

$$[u] \in K_1(B) = K_1(\mathcal{K}(V_-^{BF})) \oplus K_1(\mathcal{K}(V_+^{BF})) \cong K^1(X) \oplus K^1(X)$$

corresponds under the suspension isomorphism to

$$[\sigma_H(\tilde{P}_\gamma)] \in K_0(C_0((0,\infty),B)) \cong K_0(I_H).$$

□

In proposition 4.3.1 we have de-emphasized the Morita equivalence that is part of the definition of the map b. We specified elements

$$[\pi_\pm(\sigma_H(P_\gamma))] \in K_1(\mathcal{K}(\mathcal{H}_\pm^{BF})),$$

but we have not yet indicated what to do about the Morita equivalences

$$K_1(\mathcal{K}(\mathcal{H}_\pm^{BF})) \cong K^1(X).$$

But this is straightforward. Let $V_\pm^N \to X$ denote the complex vector bundles on X defined by

$$V_+^N = \bigoplus_{j=0}^N \operatorname{Sym}^j H^{1,0}, \ \ V_-^N = \bigoplus_{j=0}^N \operatorname{Sym}^j H^{0,1},$$

and let

$$S_\pm^N : V_\pm^{BF} \to V_\pm^N$$

be the orthogonal projection in the fibers. Let us denote by $\pi_\pm^N$ the Bargmann-Fok representations compressed by the projections S^N,

$$\pi_\pm^N(a) = S_\pm^N \pi_\pm(a) S_\pm^N.$$

For sufficiently large value of the integer N we have

$$[\pi_\pm(\sigma_H(P_\gamma))] = [S^N\pi_\pm(\sigma_H(P_\gamma))S^N + (1-S^N)] \in K_1(\mathcal{K}(\mathcal{H}_\pm^{BF})),$$

where the correct value of N depends on $\pi_\pm(\sigma_H(P_\gamma))$. Observe that with this choice of N, the elements $\pi_\pm^N(\sigma(P_\gamma))$ define automorphisms of the vector bundles $V_\pm^N$, so that

$$[V^N, \pi_\pm^N(\sigma_H(P_\gamma))] \in K^1(X).$$

The compression map

$$K_1(\mathcal{K}(\mathcal{H}_\pm^{BF})) \to K^1(X) : [\pi_\pm(\sigma_H(P_\gamma))] \mapsto [V_\pm^N, \pi_\pm^N(\sigma_H(P_\gamma))]$$

implements the Morita equivalence (where N depends on P_γ, as mentioned). So we have, in fact,

$$b(\sigma_H(P_\gamma)) = [\pi_-^N(\sigma_H(P_\gamma))] \cap [X^-] \ + \ [\pi_+^N(\sigma_H(P_\gamma))] \cap [X^+].$$

We can now prove Theorem 1.3.1 stated in the introduction, which explicitly identifies an (M,E,φ) cycle that represents $b(\sigma_H(P_\gamma))$.

Proof of Theorem 1.3.1. Because of Darboux's theorem the contact manifold can locally be identified with an open subset of the Heisenberg group $\mathbb{R}^{2n+1}$. Let X_j, Y_j, T be the standard right invariant vector fields on the Heisenberg group with $[X_j, Y_j] = T$. In such local coordinates, the operator P_γ can be then written as

$$P_\gamma = \sum_{j=1}^n -(X_j^2 + Y_j^2) + i\gamma T + \dots,$$

where we ignore first order terms in X_j, Y_j. These lower order terms may appear because the sublaplacian $\Delta_H = \sum -(X_j^2 + Y_j^2) + \dots$ is unique only up to lower order terms.

We have $JX_j = Y_j$, and so $Z_j = (X_j - iY_j)/\sqrt{2}$. A simple computation shows that $2Z_j\bar{Z}_j = X_j^2 + Y_j^2 + i[X_j, Y_j]$, and so

$$\sum -2Z_j\bar{Z}_j = \sum -(X_j^2 + Y_j^2) + inT.$$

We can therefore write

$$P_\gamma = \sum -2Z_j\bar{Z}_j - i(n-\gamma)T + \dots.$$

The Bargmann-Fok representation on $(\mathcal{H}_+^{BF})_x$ of the symbol $\sigma_H(P) = \{P_x, x \in X\}$ is

$$\pi_+(P_x) = \sum 2z_j \frac{\partial}{\partial z_j} + n - \gamma(x).$$

The symmetric powers of $H^{1,0}$ are eigenspaces of $\pi_+(P_x)$. To see this, observe that

$$\pi_+(P_x)z^\alpha = (2|\alpha| + n - \gamma(x))z^\alpha,$$

where $z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n}$ and $|\alpha| = \alpha_1 + \dots + \alpha_n$. Therefore on $\mathrm{Sym}^{\mathrm{j}} H^{1,0}$ the operator $\pi_+(P_x)$ acts as the scalar

$$a_j(x) = 2j + n - \gamma(x).$$

In the dual Bargmann-Fok representation the roles of Z, $\bar{Z}$ are reversed. So from

$$\sum -2\bar{Z}_j Z_j = \sum -(X_j^2 + Y_j^2) - inT,$$

we get

$$P_\gamma = \sum -2\bar{Z}_j Z_j + i(n+\gamma)T.$$

The representation of $\sigma_H(P)$ on the conjugate Bargmann-Fok spaces $(\mathcal{H}_-^{BF})_x$ is then given by

$$\pi_-(P_x) = \sum 2\bar{z}_j \frac{\partial}{\partial \bar{z}_j} + n + \gamma(x).$$

and we find that $\pi_-(P_x)$ acts on $\mathrm{Sym}^{\mathrm{j}} H^{0,1}$ as the scalar

$$b_j = 2j + n + \gamma(x).$$

Replacing P with

$$\tilde{P} = P(1 + P^*P)^{-1/2}$$

we obtain an order zero operator whose principal symbols act on the symmetric powers of $H^{1,0}$ by the scalar $a_j/\sqrt{1+|a_j|^2}$, and similarly for $H^{0,1}$. For large values of j these renormalized scalars are close to 1, and so they contribute only trivial summands to the K-cycle and can be ignored. We derive

$$b(\sigma_H(\tilde{P})) = \sum_j (\mathrm{Sym}^j H^{1,0}, a_j) \cap [X^+] + \sum_j (\mathrm{Sym}^j H^{0,1}, b_j) \cap [X^-].$$

We can use the non-normalized scalars, because as vector bundle automorphisms they are homotopic to their normalized versions.

□

Corollary 4.3.2. *The Chern character of the K-cycle $[P_\gamma]$ is the Poincaré dual of the cohomology class*

$$\mathrm{P.D.}\,\mathrm{ch}\,([P_\gamma]) = \sum_{j=0}^{N} \mathrm{ch}(2j+n-\gamma) \cup \mathrm{ch}(\mathrm{Sym}^{\mathrm{j}}\, H^{1,0}) \cup e^{c_1(H^{1,0})/2} \cup \hat{A}(X)$$

$$+ (-1)^{n+1} \sum_{j=0}^{N} \mathrm{ch}(2j+n+\gamma) \cup \mathrm{ch}(\mathrm{Sym}^{\mathrm{j}}\, H^{0,1}) \cup e^{c_1(H^{0,1})/2} \cup \hat{A}(X).$$

Here the odd Chern character $\mathrm{ch}(f) \in H^1(X,\mathbb{Z}) = [X, S^1]$ *refers to the* 1*-cocycle associated to a continuous map* $f : X \to \mathbb{C}^\times \sim S^1$.

Example 4.3.3. In [26] we derived an explicit geometric formula for the Fredholm index of P_γ in the simplest possible case where X is a three-dimensional manifold. Let us describe the corresponding K-cycle in this case, and compare it with the index formula in [26].

On a three-manifold X we have $\hat{A}(X) = 1$, while the bundle $H^{1,0}$ is a line bundle. Then $\mathrm{Sym}^{\mathrm{j}} H^{1,0} = (H^{1,0})^{\otimes j}$. Writing $c_1 = c_1(H^{1,0})$ we have

$$\mathrm{ch}(\mathrm{Sym}^{\mathrm{j}} H^{1,0}) e^{c_1/2} = (1 + jc_1)(1 + \frac{1}{2}c_1) = 1 + \frac{2j+1}{2}c_1.$$

Denote by

$$W_k = \mathrm{ch}(\gamma - k) = \left[-\frac{1}{2\pi i} \frac{d\gamma}{\gamma - k} \right] \in H^1(X,\mathbb{Z})$$

the 1-cocycle that encodes the winding of the coefficient γ around the odd integer k,

$$\gamma : X \to \mathbb{C} \setminus \{\text{odd integers}\}.$$

We find

$$\mathrm{P.D.}\,\mathrm{ch}(P_\gamma) = \sum_{k\,\mathrm{odd}} W_k \; + \sum_{k\,\mathrm{odd}} k\, W_k \cup \frac{c_1}{2} \; \in H^1(X,\mathbb{Z}) \oplus H^3(X,\mathbb{Z}).$$

Here all cocycles are integer cocycles. The Poincaré dual of the 3-cocycle $\sum k\, W_k \cup c_1/2$ is a 0-cycle whose image under the map $H_0(X,\mathbb{Z}) \to H_0(\mathrm{pt},\mathbb{Z}) = \mathbb{Z}$ is just the Fredholm index of P_γ. This term in our K-cycle contains exactly the same information as the index formula of [26]. The Poincaré dual of the term $\sum W_k$ is a 2-cycle. If we twist P_γ by a complex vector bundle $F \to X$, then this 2-cycle will pair with $c_1(F)$, and the curvature of F will contribute to the index of $F \otimes P_\gamma$. The information contained in the term $\sum W_k$ cannot be derived or guessed from the formulas of [25, 26]. The following formula highlights the difference between the result of the present paper and the earlier formula,

$$\mathrm{Index}(F \otimes P_\gamma) = \mathrm{rank}(F) \cdot \mathrm{Index P}_\gamma + \sum_{k\,\mathrm{odd}} \int_X \mathrm{ch}(\gamma - k) \wedge c_1(F).$$

4.4. **Toeplitz operators.** The index formula for Toeplitz operators of Louis Boutet de Monvel [12] is a corollary of Theorem 3.5.1, i.e., of commutativity of the triangle

$$\begin{tikzcd}
 & K_0(C^*(T_H X)) \arrow[dl, "b"'] \arrow[dr, "\mathrm{Op_H}"] & \\
K_0(X) \arrow[rr, "\mu"] & & KK_0(C(X),\mathbb{C})
\end{tikzcd}$$

Boutet de Monvel's formula uses the Szegö projector. If the contact manifold X is the boundary of a strictly pseudoconvex complex domain, then the Szegö projector S is defined as the projection of $L^2(X)$ onto the Hardy space $H^2(X)$. Let f be a smooth map

$$f : X \to \mathrm{GL}(r, \mathbb{C}).$$

In an evident fashion, the function f defines a multiplication operator M_f on $L^2(X)^{\oplus r}$. The Toeplitz operator T_f is the compression of M_f to $H^2(X)^{\oplus r}$, i.e.,

$$T_f = S_r M_f S_r,$$

where $S_r = S^{\oplus r}$. The Toeplitz operator T_f is a Fredholm operator, and Boutet de Monvel's formula is

$$\mathrm{Index}\, T_f = \langle \mathrm{ch}(f) \cup \mathrm{Td}(X), [X] \rangle,$$

where $\mathrm{Td}(X)$ is the Todd class of the Spin^c manifold X^+, and $[f]$ is viewed as an element in $K^1(X)$.

As shown in [17], the operator

$$\tilde{T}_f = T_f + 1 - S_r$$

is an order zero operator in the Heisenberg calculus. It is immediate that $\tilde{T}_f$ is a Fredholm operator on $L^2(X)^{\oplus r}$ with $\mathrm{Index}\, \tilde{T}_f = \mathrm{Index}\, T_f$. To prove Boutet de Monvel's formula as a corollary of Theorem 3.5.1, we use the calculation in [17] of the symbol in the Heisenberg calculus of $\tilde{T}_f$. The description in [17] is tantamount to the following. Within the Bargmann-Fok space $\mathcal{H}_+^{BF}$ there is the vacuum summand $\mathbb{C} = \mathrm{Sym}^0 \mathbb{C}^n$. Hence, the Hilbert module V_+^{BF} contains the trivial line bundle $\underline{\mathbb{C}} = \mathrm{Sym}^0 H^{1,0}$. The Bargmann-Fok representations $\pi_\pm$ of the Heisenberg symbol of $\tilde{T}_f$ are

$$\begin{aligned}
&\pi_+(\sigma_H(\tilde{T}_f)) = f \text{ acting on } \underline{\mathbb{C}}^r,\\
&\pi_+(\sigma_H(\tilde{T}_f)) = 1 \text{ on the orthogonal complement of } \underline{\mathbb{C}}^r,\\
&\pi_-(\sigma_H(\tilde{T}_f)) = 1
\end{aligned}$$

Just as in the case of the operators P_γ, the equatorial symbol of $\tilde{T}_f$ is 1, and so there exists a perturbation by $C_c^\infty(T_H X)$ so that $\pi_0(\sigma_H(\tilde{T}_f)) = 1$ as a function on H^* . This implies

$$\begin{aligned}
b(\sigma_H(\tilde{T}_f)) &= [\pi_+(\sigma_H(\tilde{T}_f))] \cap [X^+] + [\pi_-(\sigma_H(\tilde{T}_f))] \cap [X^-]\\
&= [f] \cap [X^+] \in K_0^{top}(X).
\end{aligned}$$

Equivalently, the K-cycle which solves the index problem for $\tilde{T}_f$ is $(X^+ \times S^1, E_f, \varphi)$ where E_f is the vector bundle on $X \times S^1$ obtained from a clutching construction—using trivial vector bundles of fiber dimension r—with f, and $\varphi : X \times S^1 \to X$ is the projection. In particular, the formula of Boutet de Monvel is now a corollary.

4.5. **The general case: vector bundles.** Let us briefly consider what happens if P is a hypoelliptic operator that acts on sections in a complex vector bundle,

$$P : C^\infty(X, F^0) \to C^\infty(X, F^1).$$

The case of a scalar operator twisted by a (single) vector bundle F is already contained in our earlier description. But not every operator arises in this way and it is useful to briefly repeat the steps of our procedure for constructing the (M, E, φ) cycle in the general case.

The Heisenberg symbol of an operator P acting in bundles F^0, F^1 is a smooth family $\{P_x, x \in X\}$ of translation invariant operators

$$P_x : C_c^\infty(G_x, F_x^0) \to C_c^\infty(G_x, F_x^1).$$

If P is hypoelliptic then the scalar representations of P_x assemble to a function

$$\pi_0(P_x) : H_x^* \to \mathrm{Hom}(F_x^0, F_x^1)$$

and we get a globally defined partial classical symbol

$$\pi_0(P) \in C^\infty(H^*, \mathrm{Hom}(\pi^* F^0, \pi^* F^1)),$$

where $\pi^* F^j$ is the pull-back of F^j from X to H^*. This classical part of the Heisenberg symbol is 'elliptic' in the sense that it is invertible outside a compact set in H^*. Moreover, as we have seen above, as a class in $K^0(H^*)$ the partial symbol $\pi_0(P)$ is trivial. So we can choose a compactly supported perturbation of $\pi_0(P)$ so that

$$\tilde{\pi}_0(P) \in C^\infty(H^*, \mathrm{Iso}(\pi^* F^0, \pi^* F^1)).$$

(We may have to stabilize F^0, F^1.) Such a choice is always possible, but in the general case it is not canonical, and the resulting (M, E, φ) cycle will depend on the choice made here. Extending this perturbation to the full Heisenberg symbol is precisely what is involved in choosing a lift from $K_0(C^*(T_H X))$ to $K_0(I_H)$.

Now restrict the perturbed $\tilde{\pi}_0(P)$ to the zero section in H^*, and interpret the resulting section as an isomorphism of vector bundles

$$\pi_0 : F^0 \to F^1.$$

Composing P with π_0^{-1} we obtain a new operator

$$\pi_0^{-1} \circ P : C^\infty(X, F^0) \to C^\infty(X, F^0)$$

that has, of course, the same K-homology class as P. But now the equatorial symbol of $\pi_0^{-1} \circ P$ (i.e., its value at the spherical boundary $S^* H$ of H^*) is homotopic to the constant map from $S^* H$ to the identity operator on F^0. Therefore the Bargmann-Fok representations π_+ on $V_+^{BF} \otimes F^0$ and π_- on $V_-^{BF} \otimes F^0$ of the operator $\pi_0^{-1} \circ P$ determine well-defined elements in $K^1(X)$, and we can write as before

$$\mu^{-1}(P) = b(\sigma_H(P)) = \pi_+(\sigma_H(\pi_0^{-1} \circ P)) \cap [X^+] + \pi_-(\sigma_H(\pi_0^{-1} \circ P)) \cap [X^-].$$

An (M, E, φ) cycle is constructed from this formula in the standard way.

Remark 4.5.1. The non-canonical part of this formula is the choice of the isomorphism $\pi_0 : F^0 \to F^1$. This isomorphism is chosen such that its pull-back to $S^* H$ is homotopic to the equatorial symbol of P. In the examples of the operators $P_\gamma = \Delta_H + i\gamma T$ or the Toeplitz operators an obvious choice $\pi_0 = 1$ exists, but in general the choice of π_0 is not canonical.

Example 4.5.2. With X as above, let r be a positive integer and let γ be a C^∞ map from X to the $\mathbb{C}$ vector space of all $r \times r$ matrices, denoted $M(r, \mathbb{C})$,

$$\gamma : X \to M(r, \mathbb{C})$$

Then — entirely analogous to the case $r = 1$ — a differential operator P_γ is given by :

$$P_\gamma = \Delta_H \otimes I_r + iT \otimes \gamma$$
$$P_\gamma : C^\infty(X, \mathbb{C}^r) \to C^\infty(X, \mathbb{C}^r)$$

where (as usual) I_r is the $r \times r$ identity matrix. P_γ is elliptic in the Heisenberg calculus if and only if for all $x \in X$:

$$\gamma(x) - \lambda I_r \quad \text{is invertible for all } \lambda \in \{\dots, -n-4, -n-2, -n, n, n+2, n+4, \dots\}$$

As above, P_γ determines an element $[P_\gamma]$ in $KK^0(C(X), \mathbb{C})$. The K-cycle which solves the index problem for $[P_\gamma]$ is $(X^+ \times S^1, E^+, \varphi) \sqcup (X^- \times S^1, E^-, \varphi)$ where $X^+ \times S^1 \sqcup X^- \times S^1$ and φ are as above, and

$$E^+ = \bigoplus_{j=0}^{N} \left(\nu^\gamma_{2j+1} \otimes \varphi^* \mathrm{Sym}^j H^{1,0} \right)$$

$$E^- = \bigoplus_{j=0}^{N} \left(\nu^\gamma_{-(2j+1)} \otimes \varphi^* \mathrm{Sym}^j H^{0,1} \right)$$

Here ν^γ_{2j+1} (resp. $\nu^\gamma_{-(2j+1)}$) is the $\mathbb{C}$ vector bundle of fiber dimension k on $X \times S^1$ obtained by doing a clutching construction using $\gamma - (2j+1)I_k$ (resp. $\gamma + (2j+1)I_k$).

4.6. **A hypoelliptic index problem posed by R. Ponge.** We now consider an example of a non-scalar hypoelliptic operator whose index problem was posed by Raphael Ponge in [22]. The Dirac operator, the signature operator, or the (assembled) Dolbault complex are constructed from geometric data on the manifold. No arbitrary choice of coefficients—like the coefficient γ of the operator $P_\gamma = \Delta_H + i\gamma T$—is involved in their construction. In [22] Ponge constructed a second order hypoelliptic differential operator $P = Q_b - \gamma Q_b \gamma$ that is *canonically* associated to a contact structure. Some extra structure on the manifold is involved, but it is a priori evident that the class of P in K-homology only depends on the contact structure. Ponge proved that the Connes-Moscovici local index formula represents the Chern character of P in cyclic cohomology, and posed the problem of expressing this Chern character in terms of characteristic classes of the contact structure. We are now able to answer his question, and the answer turns out to be disappointing.

Let (X, H) be a closed contact manifold with contact form θ. As above, a complex structure J is chosen for H, compatible with $d\theta$. Assume that J defines a C.R. structure, i.e., the complex bundle $H^{1,0}$ is closed under brackets of vector fields. Write $\Lambda H^* \otimes \mathbb{C} = \oplus \Lambda^{p,q}$ with $\Lambda^{p,q} = \Lambda^p (H^{1,0})^* \otimes \Lambda^q (H^{0,1})^*$, $p, q = 0, 1, \dots, n$, and let

$$\bar{\partial}_b : \Lambda^{p,q} \to \Lambda^{p,q+1}$$

be the tangential Cauchy Riemann operator on the C.R. manifold X. The operator $\bar{\partial}_b + \bar{\partial}_b^*$ is not hypoelliptic (its kernel on $\Lambda^{0,0}$ is the range of the Szegö projector), and neither is its second order analog

$$Q_b = \bar{\partial}_b^* \bar{\partial}_b - \bar{\partial}_b \bar{\partial}_b^*.$$

To construct a hypoelliptic operator Ponge makes use of a specific grading γ on the bundle $\Lambda H^* \otimes \mathbb{C}$. This grading exists if the $\mathbb{R}$ bundle H is equipped with an involution that anticommutes with J. This rather restrictive assumption amounts to the existence of a globally defined Lagrangian splitting,

$$H = E \oplus F$$

where $d\theta$ vanishes when restricted to either of the $\mathbb{R}$ vector bundles E and F, while J is an isomorphism from E to F preserving the Euclidean structures. This reduces the structure group of TX to $O(n)$. With this extra structure a conjugate linear involution $v \mapsto v^c$ can be defined for vectors $v \in H_x$ by setting $v = v^c$ if $v \in E_x$ and $v^c = -v$ for $v \in F_x$. This involution preserves the two summands in the decomposition $H \otimes \mathbb{C} = H^{1,0} \oplus H^{0,1}$, and defines conjugate linear automorphisms of these bundles. In fact, we could identify $H^{1,0} \cong H^{0,1} \cong E \otimes \mathbb{C}$.

Ponge now assumes that $\dim X = 3$, i.e., $n = 1$. The involution $v \mapsto v^c$ is then used to define a (complex linear) Hodge star operation on $\Lambda H \otimes \mathbb{C}$ by $(*\alpha)^c \wedge \beta = \langle \alpha, \beta \rangle d\theta$. Then $*^2 = (-1)^{p+q+1}$ on $\Lambda^{p,q}$ and the Hodge star gives rise to a grading operator γ on $\Lambda H \otimes \mathbb{C}$ with $\gamma^2 = 1$, $\gamma^* = \gamma$,

$$\gamma = i^{(p+q)^2+1} * : \Lambda^{p,q} \to \Lambda^{1-p,1-q}.$$

The operator

$$P = Q_b - \gamma Q_b \gamma$$

is hypoelliptic. P anticommutes with the grading γ, and so P is a Fredholm operator from sections in the bundle Λ^+ of forms with $\gamma = +1$ to the bundle Λ^- of forms with $\gamma = -1$,

$$P : C^\infty(X, \Lambda^+) \to C^\infty(X, \Lambda^-).$$

The problem posed by Ponge in [22] is to compute the Chern character of the K-homology element $[P]$. We will solve this problem here, not by using the Connes-Moscovici local index formula as proposed by Ponge, but by applying our methods to show that the K-cycle $[P]$ is always zero. While this is a disappointing result it nevertheless proves the effectiveness of our approach, since no resolution of Ponge's question had been obtained before now.

Proposition 4.6.1. *The K-homology class $[P] \in KK^0(C(X), \mathbb{C})$ determined by Ponge's operator $P = Q_b - \gamma Q_b \gamma$ is zero.*

Proof. We first need to find the model operator P_x by freezing coefficients of P. Let $G = \mathbb{R}^3$ be the Heisenberg group with coordinates (x, y, t). The translation invariant differential forms on G are spanned by dx, dy, θ, where θ is the standard contact form

$$\theta = dt + \frac{1}{2}(y dx - x dy).$$

We trivialize the bundle $\Lambda^{*,*} = G \times \mathbb{C}^4$ by choosing basis vectors 1 for $\Lambda^{0,0}$, $dz = (dx + i dy)/\sqrt{2}$ for $\Lambda^{0,1}$, $d\bar{z} = (dx - i dy)/\sqrt{2}$ for $\Lambda^{1,0}$, and $d\bar{z} \wedge dz = i dx \wedge dy$ for $\Lambda^{1,1}$. The tangential C.R. operator on $\mathbb{R}^3$ is formally given by

$$\bar{\partial}_b = d\bar{z} \wedge \bar{Z} : \Lambda^{p,q} \to \Lambda^{p,q+1},$$

where $\bar{Z} = (X + iY)/\sqrt{2}$. Using the trivialization of $\Lambda^{*,*}$ and the equality $Z^* = -\bar{Z}$ we find a matrix expression for Q_b,

$$\bar{\partial}_b = \begin{pmatrix} 0 & 0 & 0 & 0 \\ \bar{Z} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \bar{Z} & 0 \end{pmatrix} \qquad Q_b = \begin{pmatrix} -Z\bar{Z} & 0 & 0 & 0 \\ 0 & \bar{Z}Z & 0 & 0 \\ 0 & 0 & -Z\bar{Z} & 0 \\ 0 & 0 & 0 & \bar{Z}Z \end{pmatrix}$$

With $\Delta_H = -X^2 - Y^2$ we have $-Z\bar{Z} = \frac{1}{2}(\Delta_H - iT)$ and $Z\bar{Z} = \frac{1}{2}(-\Delta_H - iT)$. We see that Q_b is not a Rockland operator.

To get an explicit expression for the model operator P on $G = \mathbb{R}^3$ we need to analyze the grading γ. The involution on $\Lambda^{*,*}$ preserves the standard basis (up to factors i^m), and is conjugate linear. A straightforward calculation shows that

$$\gamma(1) = d\bar{z} \wedge dz,\ \gamma(d\bar{z}) = -i dz,\ \gamma(dz) = +i d\bar{z},\ \gamma(d\bar{z} \wedge dz) = 1$$

or in matrix form

$$\gamma = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

The result is

$$P = \begin{pmatrix} \Delta_H & 0 & 0 & 0 \\ 0 & -\Delta_H & 0 & 0 \\ 0 & 0 & \Delta_H & 0 \\ 0 & 0 & 0 & -\Delta_H \end{pmatrix}$$

If we choose an appropriate basis for $\Lambda^{*,*}$ consisting of ± 1 eigenvectors for the grading γ, then the odd operator P takes the form

$$P = \begin{pmatrix} 0 & P^- \\ P^+ & 0 \end{pmatrix}, \qquad P^+ = (P^+)^* = P^- = \begin{pmatrix} \Delta_H & 0 \\ 0 & -\Delta_H \end{pmatrix}$$

On a contact manifold of dimension three the operator on $\mathbb{R}^3$ that we computed here describes the model operator P_x at each point $x \in X$. The equatorial symbol of P_x is *constant* along the spheres S^*H_x for fixed $x \in X$, and so the lift of the symbol $\sigma_H(P)$ to $K_0(I_H)$ is canonical and unproblematic. In the matrix notation above

$$\pi_0(P) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

with the appropriate choice of basis for $\Lambda^{\pm}$. We obtain an isomorphism of vector bundles $\pi_0 : \Lambda^+ \to \Lambda^-$. If we compose P with π_0 then the modified operator $\pi_0 \circ P$ has symbol

$$\pi_0 \circ P_x = \begin{pmatrix} \Delta_H & 0 \\ 0 & \Delta_H \end{pmatrix}$$

This means that the Heisenberg symbol of the operator $\pi_0 \circ P$ is the symbol of a sublaplacian Δ_H on X twisted by the vector bundle Λ^+. Therefore in K-homology $[P] = [\pi_0 \circ P]$ equals $[\Delta_H]$ twisted by the vector bundle Λ^+. Since the element in K-homology $[\Delta_H] = 0$ it follows that $[P] = 0$.

□

Remark 4.6.2. Although somewhat disappointing, the topological triviality of the Ponge operator can be understood as an example of a general phenomenon. Canonical geometric operators, e.g., Dirac operator, signature operator, Dolbeault operator, Szegö projector, are associated to a reduction of structure group of the tangent bundle TX of a manifold. Let G denote a compact Lie group acting on a real vector space V, such that $TX \cong P \times_H V$, where P is a principal G-bundle over X. As discussed by Atiyah and Singer [3, section 2], there is a natural homomorphism

$$\alpha_P : K^0_G(V^*) \to K^0(T^*X)$$

which maps a 'universal' elliptic symbol class $v \in K^0_G(V^*)$ in the G-equivariant K-theory group of V^* to an elliptic symbol class in $K^0(T^*X)$ associated to the G-structure. All (symbols of) classical geometric operators arise in this way.

This construction can be adapted to the Heisenberg calculus as follows. Let $V = \mathbb{C}^n \times \mathbb{R}$ be the Heisenberg group, and G a compact Lie group that acts on V by group automorphisms. Then the bundle of osculating groups T_HX associated to a contact structure on X is an associated bundle $T_HX \cong P \times_G V$, where P is a principal G-bundle over X. For example, a choice of complex structure J for the contact hyperplane bundle $H \subset TX$ reduces the structure group to $G = U(n)$. In Ponge's example it is further reduced to $G = O(n)$.

A canonical geometric operator in the Heisenberg calculus (or rather its symbol) then arises as an element in the range of a natural homomorphism

$$\alpha_P : K_0^G(C^*(V)) \to K_0(C^*(T_H X)).$$

But V is odd dimensional, and the group $K_0^G(C^*(V)) \cong K_G^0(V^*)$ is zero. Therefore the image in the even K-homology group

$$\mathrm{Op}_H : K_0(C^*(T_H X)) \cong KK^0(C(X), \mathbb{C})$$

is automatically zero for *any* canonical geometric hypoelliptic operator in the Heisenberg calculus.

To get nonzero elements in the even K-homology group a canonical geometric operator must be paired with a K^1 element, and thus the resulting operator is not canonical since the choice of K^1 element is arbitrary. Examples of this are the Toeplitz operators T_f and the operators P_γ considered above. The construction of the Ponge operator $P = Q_b - \gamma Q_b \gamma$ is canonical (i.e., has no input from K^1) and thus it is not surprising that it is zero in even K-homology.

4.7. **Odd-dimensional spheres.** S^{2n+1}, as usual, is the unit sphere of $\mathbb{R}^{2n+2}$. In this section we show how the general theory developed above applies to give a very simple and clear index formula for a naturally arising class of hypoelliptic operators on S^{2n+1}. The standard contact form θ for S^{2n+1} is

$$\theta = \sum_{i=1}^{n+1} x_{2i} dx_{2i-1} - x_{2i-1} dx_{2i}$$

where $x_1, x_2, \ldots, x_{2n+2}$ are the standard coordinates on $\mathbb{R}^{2n+2}$. Let r be a positive integer, and let $\gamma : S^{2n+1} \longrightarrow M(r, \mathbb{C})$ be a smooth map from S^{2n+1} to $M(r, \mathbb{C})$ where $M(r, \mathbb{C})$ is the $\mathbb{C}$ vector space of all $r \times r$ complex matrices. We require that for all $p \in S^{2n+1}$

$$\gamma(p) - \lambda I \quad \text{is invertible for all } \lambda \in \{\ldots, -n-4, -n-2, -n, n, n+2, n+4, \ldots\}$$

Associated to γ is a Fredholm operator P_γ

$$P_\gamma : C^\infty(S^{2n+1}, \mathbb{C}^r) \longrightarrow C^\infty(S^{2n+1}, \mathbb{C}^r)$$

defined by

$$P_\gamma = \Delta_H \otimes I_r \; + \; iT \otimes \gamma$$

where, as above, Δ_H is the sub-Laplacian and T is the Reeb vector field,

$$T = \sum_{i=1}^{n+1} x_{2i} \frac{\partial}{\partial x_{2i-1}} - x_{2i-1} \frac{\partial}{\partial x_{2i}}$$

Any continuous map $f : S^{2n+1} \longrightarrow GL(r, \mathbb{C})$ determines an element in $K^1(S^{2n+1})$. By Bott periodicity, $K^1(S^{2n+1}) = \mathbb{Z}$. Therefore an integer $\beta(f)$ has been assigned to f,

$$\beta(f) = ch(f)[S^{2n+1}] = \int_{S^{2n+1}} \mathrm{Tr} \left(\frac{f^{-1} df}{-2\pi i} \right)^{2n+1}$$

The formula for the index of P_γ is

$$\mathrm{Index}\, P_\gamma = \sum_{j=0}^{N} \binom{n+j-1}{j} \left[\beta(\gamma - (n+2j) I_r) + (-1)^{n+1} \beta(\gamma + (n+2j) I_r) \right]$$

where N is sufficiently large, as above. This formula is obtained by applying example 4.5.2. Observe that on an odd dimensional sphere any complex vector bundle is stably trivial. Thus $H^{1,0}$ is stably trivial, so that in the K-cycle of example 4.5.2 we may replace $\varphi^*\mathrm{Sym}^j H^{1,0}$ by a trivial vector bundle of the same rank, which is $\begin{pmatrix} n+j-1 \\ j \end{pmatrix}$.

In particular, on S^3 the formula is

$$\mathrm{Index}\, P_\gamma = \sum_{j=0}^{N} \beta(\gamma - (2j+1)I_r) + \beta(\gamma + (2j+1)I_r) = \sum_{k\,\mathrm{odd}} \beta(\gamma - kI_r)$$

References

[1] M. F. Atiyah, *Global theory of elliptic operators*, Proc. Internat. Conf. on Functional Analysis and Related Topics (Tokyo, 1969), 1970, pp. 21–30.

[2] M. F. Atiyah and I. M. Singer, *The index of elliptic operators. I*, Ann. of Math. (2) **87** (1968), 484–530.

[3] ______, *The index of elliptic operators. III*, Ann. of Math. (2) **87** (1968), 546–604.

[4] V. Bargmann, *On a Hilbert space of analytic functions and an associated integral transform*, Comm. Pure Appl. Math. **14** (1961), 187–214.

[5] P. Baum and R. G. Douglas, *Index theory, bordism, and K-homology*, Operator algebras and K-theory (San Francisco, Calif., 1981), 1982, pp. 1–31.

[6] ______, *K homology and index theory*, Operator algebras and applications, Part I (Kingston, Ont., 1980), 1982, pp. 117–173.

[7] P. Baum, N. Higson, and T. Schick, *On the equivalence of geometric and analytic K-homology*, Pure Appl. Math. Q. **3** (2007), no. 1, part 3, 1–24.

[8] P. Baum and E. van Erp, *Conjugation in K-homology*, forthcoming.

[9] ______, *K-homology and the index of elliptic operators*, forthcoming.

[10] R. Beals and P. Greiner, *Calculus on Heisenberg manifolds*, Annals of Mathematics Studies, vol. 119, Princeton University Press, Princeton, NJ, 1988.

[11] B. Blackadar, *K-theory for operator algebras*, Second, Mathematical Sciences Research Institute Publications, vol. 5, Cambridge University Press, Cambridge, 1998.

[12] L. Boutet de Monvel, *On the index of Toeplitz operators of several complex variables*, Invent. Math. **50** (1978/79), no. 3, 249–272.

[13] M. D. Choi and E. G. Effros, *The completely positive lifting problem for C^*-algebras*, Ann. of Math. (2) **104** (1976), no. 3, 585–609.

[14] A. Connes and H. Moscovici, *The local index formula in noncommutative geometry*, Geom. Funct. Anal. **5** (1995), no. 2, 174–243.

[15] ______, *Hopf algebras, cyclic cohomology and the transverse index theorem*, Comm. Math. Phys. **198** (1998), no. 1, 199–246.

[16] A. Connes, *Noncommutative geometry*, Academic Press Inc., San Diego, CA, 1994.

[17] C. L. Epstein, *Lectures on indices and relative indices on contact and CR-manifolds*, Woods Hole mathematics, 2004, pp. 27–93.

[18] G. B. Folland and E. M. Stein, *Estimates for the $\bar\partial_b$ complex and analysis on the Heisenberg group*, Comm. Pure Appl. Math. **27** (1974), 429–522.

[19] M. Hilsum and G. Skandalis, *Morphismes K-orientés d'espaces de feuilles et fonctorialité en théorie de Kasparov (d'après une conjecture d'A. Connes)*, Ann. Sci. École Norm. Sup. (4) **20** (1987), no. 3, 325–390.

[20] R. Melrose, *Homology and the Heisenberg algebra*, Séminaire sur les Équations aux Dérivées Partielles, 1996–1997, 1997, pp. Exp. No. XII, 11. Joint work with C. Epstein and G. Mendoza.

[21] V. Nistor, *An index theorem for gauge-invariant families: the case of solvable groups*, Acta Math. Hungar. **99** (2003), no. 1-2, 155–183.

[22] R. Ponge, *Géométrie spectrale et formules d'indices locales pour les variétés CR et contact*, C. R. Acad. Sci. Paris Sér. I Math. **332** (2001), no. 8, 735–738.

[23] M. E. Taylor, *Noncommutative microlocal analysis. I*, Mem. Amer. Math. Soc. **52** (1984), no. 313, iv+182.

[24] E. van Erp, *The Atiyah-Singer formula for subelliptic operators on a contact manifold, Part I*, Ann. of Math. **171** (2010), 1647–1681.

[25] ______, *The Atiyah-Singer formula for subelliptic operators on a contact manifold, Part II*, Ann. of Math. **171** (2010), 1683–1706.

[26] ———, *Noncommutative topology and the world's simplest index theorem*, Proc. Natl. Acad. Sci. USA **107** (2010), no. 19, 8549–8556.
[27] ———, *The index of hypoelliptic operators on foliated manifolds*, J. Noncommut. Geom. (2011), in press.

The Pennsylvania State University, University Park, PA, 16802, USA
E-mail address: `baum@math.psu.edu`

Dartmouth College, 6188, Kemeny Hall, Hanover, New Hampshire, 03755, USA
Current address: University of Hawaii at Manoa, 2565 McCarthy Mall, Honolulu, Hawaii 96822, USA
E-mail address: `jhamvanerp@gmail.com`